



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2011

---

## **Adaptation of *Saccharomyces cerevisiae* to saline stress through laboratory evolution**

Dhar, R ; Sägesser, R ; Weikert, C ; Yuan, J ; Wagner, A

**Abstract:** Most laboratory evolution studies that characterize evolutionary adaptation genomically focus on genetically simple traits that can be altered by one or few mutations. Such traits are important, but they are few compared with complex, polygenic traits influenced by many genes. We know much less about complex traits, and about the changes that occur in the genome and in gene expression during their evolutionary adaptation. Salt stress tolerance is such a trait. It is especially attractive for evolutionary studies, because the physiological response to salt stress is well-characterized on the molecular and transcriptome level. This provides a unique opportunity to compare evolutionary adaptation and physiological adaptation to salt stress. The yeast *Saccharomyces cerevisiae* is a good model system to study salt stress tolerance, because it contains several highly conserved pathways that mediate the salt stress response. We evolved three replicate lines of yeast under continuous salt (NaCl) stress for 300 generations. All three lines evolved faster growth rate in high salt conditions than their ancestor. In these lines, we studied gene expression changes through microarray analysis and genetic changes through next generation population sequencing. We found two principal kinds of gene expression changes, changes in basal expression (82 genes) and changes in regulation (62 genes). The genes that change their expression involve several well-known physiological stress-response genes, including CTT1, MSN4 and HLR1. Next generation sequencing revealed only one high-frequency single-nucleotide change, in the gene MOT2, that caused increased fitness when introduced into the ancestral strain. Analysis of DNA content per cell revealed ploidy increases in all the three lines. Our observations suggest that evolutionary adaptation of yeast to salt stress is associated with genome size increase and modest expression changes in several genes.

DOI: <https://doi.org/10.1111/j.1420-9101.2011.02249.x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-59820>

Journal Article

Accepted Version

Originally published at:

Dhar, R; Sägesser, R; Weikert, C; Yuan, J; Wagner, A (2011). Adaptation of *Saccharomyces cerevisiae* to saline stress through laboratory evolution. *Journal of Evolutionary Biology*, 24(5):1135-1153.

DOI: <https://doi.org/10.1111/j.1420-9101.2011.02249.x>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

**Title: Adaptation of *Saccharomyces cerevisiae* to saline stress through laboratory evolution**

**Riddhiman Dhar<sup>1,3</sup>, Rudolf Sägesser<sup>1,3</sup>, Christian Weikert<sup>1,3</sup>, Ju Yuan<sup>1,3</sup> and Andreas Wagner<sup>1,2,3,4</sup>**

<sup>1</sup>Dept. of Evolutionary Biology and Environmental Studies, University of Zurich

<sup>2</sup> The Santa Fe Institute

<sup>3</sup> The Swiss Institute of Bioinformatics

<sup>4</sup> Author for correspondence

Address:  
University of Zurich  
Dept. of Biochemistry, Bldg. Y27  
Winterthurerstrasse 190  
CH-8057 Zurich  
Switzerland

Email: [aw@bioc.uzh.ch](mailto:aw@bioc.uzh.ch)  
Phone: +41-44-635-6141  
Fax: +41-44-635-6144

Formatiert: Französisch (Schweiz)

**Abstract**

Most laboratory evolution studies that characterize evolutionary adaptation genomically focus on genetically simple traits that can be altered by one or few mutations. Such traits are important, but they are few compared to complex, polygenic traits influenced by many genes. We know much less about complex traits, and about the changes that occur in the genome and in gene expression during their evolutionary adaptation. Salt stress tolerance is such a trait. It is especially attractive for evolutionary studies, because the physiological response to salt stress is well-characterized on the molecular and transcriptome level. This provides a unique opportunity to compare evolutionary adaptation and physiological adaptation to salt stress. The yeast *Saccharomyces*

*cerevisiae* is a good model system to study salt stress tolerance, because it contains several highly conserved pathways that mediate the salt stress response. We evolved three replicate lines of yeast under continuous salt (NaCl) stress for 300 generations. All three lines evolved faster growth rate in high salt conditions than their ancestor. In these lines, we studied gene expression changes through microarray analysis, and genetic changes through next generation population sequencing. We found two principal kinds of gene expression changes, changes in basal expression (82 genes), and changes in regulation (62 genes). The genes that change their expression involve several well-known physiological stress response genes, including *CTT1*, *MSN4*, *HLR1*. Next generation sequencing revealed only one high frequency single nucleotide change, in the gene *MOT2*, that caused increased fitness when introduced into the ancestral strain. Analysis of DNA content per cell revealed ploidy increases in all the three lines. Our observations suggest that *evolutionary* adaptation of yeast to salt stress is associated with genome size increase and modest expression changes in several genes.

59  
60  
61  
62  
63  
64  
65  
66  
67  
68

## 69 Introduction

The ability to characterize the changes that occur during evolutionary adaptation on a genome-wide scale has been a boon for the field of laboratory evolution. Most published studies focus on traits with a simple basis, where changes of major effects in one or few changes can alter a trait during laboratory evolution experiments (Ferea *et al.*, 1999; Blanc & Adams, 2003; Velicer *et al.*, 2006; Stanek *et al.*, 2009). Such traits are important, but they are in the minority. The vast majority of traits have a complex, polygenic basis (refs. Benfey, *Genomics textbook*). We know much less about how genomic change and change in gene expression occurs in such polygenic traits. Our study is a step towards answering this question. We here focus on a prototypical polygenic trait, an organism's response to high concentrations of salt in its environment. The physiological response of an organism to such salt stress is well-studied on the molecular and the transcriptome levels (Posas *et al.*, 2000; Rep *et al.*, 2000; Causton *et al.*, 2001; Yale & Bohnert, 2001). This fact provides another important motivation to study the salt stress response in an evolutionary context. It allows us to ask whether evolutionary adaptation to salt stress is similar to physiological adaptation, using a genome-scale approach that relies on transcriptome changes in response to salt stress. Environmental fluctuations and stressors constantly challenge organisms in the wild. Organisms thus use cellular mechanisms to adapt to and to survive environmental fluctuations. Hyperosmotic stress is one prominent environmental

89 stressor, where a cell experiences higher solute concentration outside the cell than  
90 inside. This causes water loss from the cell, resulting in a higher intracellular  
91 concentration of ions and metabolites, and eventual arrest of cellular activity.  
92 Hyperosmotic stress is caused by high concentrations of sugar or salt, e.g., sodium  
93 chloride (NaCl). High salt stress is a special case of hyperosmotic stress and has similar  
94 effects on a cell as high concentration of sugars (Gasch *et al.*, 2000; Causton *et al.*,  
95 2001). In addition, it causes hyperionic stress due to high extracellular concentrations of  
96 Na<sup>+</sup> and Cl<sup>-</sup> ions, which are imported into the cell and can disrupt cellular ionic  
97 equilibrium. Tolerance to Na<sup>+</sup> stress thus needs additional ion transport and  
98 detoxification mechanisms along with those required for the hyperosmotic stress  
99 response (Serrano, 1996; Apse *et al.*, 1999; Gaxiola *et al.*, 1999; Maathuis & Amtmann,  
100 1999; Serrano *et al.*, 1999; Zhu, 2001). The hyperosmotic stress response in yeast is  
101 mediated by the high osmolarity glycerol (HOG) pathway, which is a MAPK pathway  
102 (Brewster *et al.*, 1993; Dihazi *et al.*, 2004; Saito & Tatebayashi, 2004; reviewed in  
103 Hohmann 2002). Two cell membrane-bound sensors Sho1p and Sln1p detect osmotic  
104 change, which results in activation of Hog pathway genes, which, in turn, leads to the  
105 activation of the downstream genes associated with salt tolerance and adaptation  
106 (Martinez-Pastor *et al.*, 1996; Schmitt & McEntee, 1996; Gorner *et al.*, 1998; Ostrander  
107 & Gorman, 1999; Rep *et al.*, 1999; Reiser *et al.*, 2000; Rep *et al.*, 2000). Whole  
108 transcriptome studies by Gasch *et al.*, Posas *et al.*, Yale and Bonhart, Causton *et al.*, and  
109 Rep *et al.* (Gasch *et al.*, 2000; Posas *et al.*, 2000; Rep *et al.*, 2000; Causton *et al.*, 2001;  
110 Yale & Bohnert, 2001) have identified hundreds of genes whose expression levels are  
111 affected by hyperosmotic stress. The genes induced during osmotic stress response  
112 include the genes involved in synthesis and regulation of the cellular osmolytes glycerol  
113 and trehalose. Another group of genes, specifically activated under saline stress, are  
114 associated with ion homeostasis. Some of the genes affected by osmotic stress show  
115 transient expression changes, whereas others show expression changes that are stable  
116 in time. In addition to genome wide approaches, many studies have characterized the  
117 roles of individual genes in the osmotic stress response of yeast (Haro *et al.*, 1991;  
118 Garciadeblas *et al.*, 1993; Mai & Breeden, 1997; Ganster *et al.*, 1998; Mendizabal *et al.*,  
119 1998; Tsujimoto *et al.*, 2000; Betz *et al.*, 2002; Goossens *et al.*, 2002; Hirata *et al.*,  
120 2003; Heath *et al.*, 2004; ).

121 Yeast is a good model system for studying osmo-adaptation in eukaryotes,  
122 since other fungi and plants share many of the stress response pathways and proteins  
123 involved in osmo-adaptation in yeast. First, the mitogen-activated protein kinase (MAP  
124 kinase) cascade, central to the stress response in yeast, is a conserved eukaryotic  
125 signal transduction pathway present from fungi to plants (Pelech & Sanghera, 1992;  
126 Errede & Levin, 1993; Nishida & Gotoh, 1993; Neiman, 1993; Neiman *et al.*, 1993;  
127 Kosako *et al.*, 1993; Tsuda *et al.*, 1993; Yashar *et al.*, 1993; Cooper, 1994; Zamanillo *et al.*,  
128 1994; Sanchez *et al.*, 1994; Yan *et al.*, 1994; Galcheva-Gargova *et al.*, 1994; Han *et al.*,  
129 1994; Waskiewicz & Cooper, 1995; Hirt, 1997; Jonak *et al.*, 1999; Kovtun *et al.*,  
130 2000; Kyriakis & Avruch, 2001). Stress signaling in plants is also carried out by MAPK  
131 pathways, which are activated by cold, drought, salt, heat, and oxidative stress (Jonak  
132 *et al.*, 1996; Kovtun *et al.*, 2000; Teige *et al.*, 2004). Second, yeast and plants have

highly similar genes required for stress tolerance (Mendoza *et al.*, 1994; Bressan *et al.*, 1998; Pardo *et al.*, 1998; Lee *et al.*, 1999; Sanders, 2000; Hasegawa & Bressan, 2000; Quintero *et al.*, 2002; Zhu, 2002). Third, yeast and plants have similar membrane ion transport and detoxification systems (Gaxiola *et al.*, 1999; Quintero *et al.*, 2000). For example, HAL family genes are important for ion homeostasis in yeast as well as in plants (Haro *et al.*, 1991; Gaxiola *et al.*, 1992; Ferrando *et al.*, 1995; Murguía *et al.*, 1996; Rios *et al.*, 1997; Mulet *et al.*, 1999; Espinosa-Ruiz *et al.*, 1999; Gisbert *et al.*, 2000; Yang *et al.*, 2001).

With one exception (Samani & Bell, 2010), all previous studies of salt stress adaptation in yeast focused on *physiological* adaptation. Such adaptation occurs on time scales up to a few hours. The mechanisms of longer term *evolutionary* adaptation to salt stress are not known. Such adaptations occur on time scales of hundred generations or more. One aim of this study is to compare a population's evolutionary response to salt stress with its physiological response on the transcriptome level. Does evolutionary adaptation mirror physiological adaptation? Does it affect largely the same genes as the physiological response?

A second aim is to investigate the genetic basis of *evolutionary* adaptation of yeast to high saline stress, as far as this is possible for a complex trait. Does the adaptation come about through accumulation of identifiable beneficial point mutations? Does it involve chromosomal rearrangements as observed in evolution of yeast in glucose-limited or phosphate-limited media? Or does it take place through genetic and epigenetic changes altering expression of genes that help cells adapt to high salt? These are some of the questions we ask.

Understanding of *evolutionary* principles of salt tolerance could also be important for biotechnological applications. Firstly, yeast cells experience high salt concentration in many industrial fermentation processes, and improvement in performance of yeast in such conditions would benefit the industry immensely (Attfield, 1997; Trainotti & Stambuk, 2001; Zheng *et al.*, 2011). Secondly, the principles of salt tolerance in yeast might be useful for engineering fungi and crop plants for salt tolerance, as both classes of organisms share many components of their stress response systems.

In this contribution, we exposed yeast to high salt concentrations for 300 generations in the laboratory in medium containing sodium chloride (NaCl). We compared the fitness and viability of the evolved lines with the starting yeast strain (ancestral strain), followed by characterization of gene expression changes and genetic changes such as mutations, copy number variations and chromosomal alterations.

## Results

To test for evolutionary adaptation of yeast cells to osmotic stress, we carried out laboratory evolution experiments through 30 serial transfers in batch cultures, comprising approximately 300 cell generations (see Methods). This number of generations is sufficient to show evolutionary adaptations in yeast (Adams & Oeller, 1986; Adams *et al.*, 1992; Dunham *et al.*, 2002; Gresham *et al.*, 2008). For our experiments, we used haploid populations to avoid any potential masking of adaptively

177 significant alleles in the diploid stage (Zeyl *et al.*, 2003). We carried out three parallel  
178 replicate evolution lines in which yeast cells grew and divided in yeast peptone medium  
179 supplemented with galactose as the sole carbon source and with 0.5M NaCl (YPGN),  
180 which is a high salt medium that exposes cells to high osmolarity stress. We refer to  
181 these lines as S lines (S1, S2 and S3). As a control, we also carried out three replicates  
182 where the growth medium did not contain NaCl (lines C1, C2 and C3).

### 184 **Evolutionary adaptation to NaCl**

185 Before embarking on our experiments, we asked whether NaCl affects the fitness of our  
186 ancestral strain. Only in this case would we expect that NaCl exerts a selection  
187 pressure to which the strain can adapt. Fitness has two main components in our  
188 experiments. These are viability on the one hand, and growth or cell division rate on the  
189 other hand. We found that osmotic stress does not affect viability of the ancestral strain  
190 significantly (Figure 1a). Not surprisingly then, viability does not increase over the  
191 course of our experiment for lines evolved on NaCl (Figure 1b) and for lines evolved  
192 without NaCl (Figure 1c).

193 In contrast to viability, population growth rate is affected by salt stress, as  
194 shown in supplementary figure 1. The figure indicates that the ancestral strain grows  
195 significantly more slowly in medium containing NaCl. **Thus, the fitness of the ancestral**  
196 **strain is lower in salt compared to that of the ancestral strain in medium without any salt.**  
197 Having established that NaCl does affect the fitness of the ancestral strain through its  
198 growth rate, we asked if the growth rate and thus, relative fitness  $w$  (see Methods) of  
199 three salt evolved lines have increased after 300 generations of evolution (Figure 2a).  
200 Fitness increased significantly relative to the ancestral strain, such that the final evolved  
201 lines had a relative population mean fitness between  $w=1.11$  and  $w=1.17$ , which  
202 corresponds to a decrease in the average cell doubling time between 8.2% and 12.3%  
203 relative to the ancestral strain (see Methods).

204 Since our high salt medium is a complex medium, it is likely that part of the  
205 evolutionary response we observe also reflects adaptation to medium components  
206 different from NaCl. To ask whether this is the case, we also measured the relative  
207 fitness of the three S lines in the control medium (without NaCl). Not surprisingly, the  
208 fitness in this medium had also increased (Figure 2b) relative to the ancestral strain,  
209 which suggests that at least part of the evolutionary adaptation we see is not specific to  
210 NaCl as a stressor. However, two lines of evidence show that a substantial fraction of  
211 the fitness increase is specific to NaCl. The first is that the fitness increase, when  
212 measured in the absence of NaCl, appears much lower than when measured in the  
213 presence of NaCl (Figures 2a and 2b) (**t-test,  $p<0.0001$  for S1, S2 and S3**). The second  
214 line of evidence is provided by our three control lines that had evolved without the  
215 addition of NaCl to the medium. Figure 2a shows that in every single line the fitness  
216 increase, when measured in medium with NaCl, is consistently and significantly lower  
217 than for the lines that had evolved in NaCl (**t-test,  $p<0.0001$  for comparisons between**  
218 **S1 & C1, S2 & C2 and S3 & C3**). Specifically, the increase in relative fitness  $w$  was at  
219 least 54 percent higher in the lines evolved in NaCl compared to the lines evolved  
220 without NaCl. In sum, a significant and substantial part of the evolutionary adaptation

we observe is due to the selection pressure provided by NaCl.

### Gene Expression analysis of the evolved lines

To compare the expression levels of genes in the evolved lines and in the ancestral strain, we performed whole gene transcriptome analysis using yeast microarrays (Affymetrix). Because previous experiments had shown that many yeast genes change expression in response to salt and to other stressors (Gasch *et al.*, 2000; Posas *et al.*, 2000; Rep *et al.*, 2000; Causton *et al.*, 2001; Yale & Bohnert, 2001), we expected that this would also hold for our lines. Thus, we first asked which genes respond to salt in a similar manner in the ancestral strain and in all evolved lines. We will refer to such genes for brevity as *shared* genes. **For all the analyses presented hereon we used all the replicates (see methods).**

In all, we observed 581 shared genes that were induced in salt and 580 shared genes that were repressed in salt with 2-fold expression change ( $|\log_2(\text{fold change})| \geq 1$ , t-test, p-value < 0.05, False Discovery Rate < 10%). Figure 3a shows a “volcano plot” ( $-\log_{10}(\text{p-value})$  vs  $\log_2(\text{fold change})$ ) for all the yeast genes in response to NaCl in the S lines and the ancestral strain. Figure 3b shows the extent of expression change in the evolved lines (vertical axis) and in the ancestral strain (horizontal axis) for all genes with similar response to NaCl in the evolved lines and the ancestral strain ( $|\log_2(\text{fold change})| \geq 1$ , t-test, p-value < 0.05, False Discovery Rate < 10%). The genes we labeled by name in the figure include some known stress response genes (e.g., *GPD1*, *SIP18*), some genes whose expression changed substantially (e.g., *FMP48*, *NOG1*), and genes in both categories (e.g., *GRE1*). Many of these shared genes are directly associated with saline stress, hyperosmotic stress, or the general stress response, and were shown to be affected by hyper-osmolarity and salt stress in previous studies of the *physiological* and osmotic stress response (Gasch *et al.*, 2000; Posas *et al.*, 2000; Rep *et al.*, 2000; Causton *et al.*, 2001; Yale & Bohnert, 2001). Among the 581 shared induced genes, 192 genes were shown to be affected by stress in previous studies; among the 580 shared repressed genes, 56 were affected by stress in previous studies.

Next, we classified shared genes using the Comprehensive Yeast Genome Database (CYGD) classification from the Munich Information Center for Protein Sequences (MIPS) (Guldener *et al.*, 2005). A detailed analysis is given in the supplementary material (supplementary figure 2 and supplementary text). Here, we only discuss two gene classes in more detail. First, genes associated with 'cell rescue, defense and virulence' contain many general stress response genes, as well as genes that respond specifically to saline stress. One would expect that such genes are induced in response to salt stress, and our data show that induced genes in this category are overrepresented, and repressed genes in this category are underrepresented (supplementary figure 2 and supplementary text). This group of genes was also found to be significantly overrepresented among genes whose deletion reduced the growth rate of yeast in salt (Warringer *et al.*, 2003). A second class of genes are genes involved in protein synthesis. The *physiological* response to stress can cause repression of protein synthesis (Gasch *et al.*, 2000). In support of this observation, our data show that significantly fewer genes associated with protein

265 synthesis are induced, and significantly more genes than expected are repressed  
266 (supplementary figure 2 and supplementary text). Again, this group of genes was  
267 significantly overrepresented among genes whose deletion increased salt-resistance  
268 (Warringer *et al.*, 2003).

269

## 270 **Differentially Expressed Genes**

271 While an analysis of genes regulated *similarly* in ancestral and evolved  
272 strains is instructive, we were more interested in genes that are expressed *differently*.  
273 These are genes whose expression adapted evolutionarily to salt stress. They can be  
274 subdivided into two main categories. The first comprises genes whose regulation has  
275 changed in the evolved lines. Figure 4a and b show two hypothetical examples of genes  
276 in this category. The second category comprises genes whose basal level of expression  
277 changed in the evolved lines, even in the absence of salt stress (Figure 4c and d). This  
278 second category of genes may be especially important, because our selection  
279 conditions imposed continuous salt stress. It is thus conceivable that cells whose  
280 expression is ancestrally regulated in response to salt stress simply increase or  
281 decrease their basal expression to the level of regulated expression in the ancestral  
282 strain in the absence of salt. We note that these two categories have multiple  
283 subcategories, and there can be genes where both the basal expression and regulation  
284 can change. Supplementary figure 3 shows an overview of all possibilities.

285 We distinguished genes in the main categories by calculating two different Z-  
286 scores for each gene, a Z-score for change in basal expression ( $Z_b$ ) and a Z-score for  
287 change in regulation ( $Z_r$ ) (see Methods) (Mukhopadhyay *et al.*, 2006). In this analysis,  
288 we considered genes with  $|Z\text{-score}| \geq 1.5$  as differentially expressed.

289

## 290 **Genes whose regulation changes**

291 In all, there are 62 genes whose regulation changes during our experiment  
292 (Figure 5a). As we mentioned above, multiple types of such change are possible.  
293 Firstly, a gene's induction in salt can increase. This will occur if the gene is induced  
294 more strongly in the evolved lines than the ancestral strain (supplementary figure 3b).  
295 Secondly, a gene's induction in salt can become reduced (supplementary figure 3c).  
296 Thirdly, a gene's repression in salt can be increased, i.e., the gene becomes repressed  
297 to a greater extent in the evolved lines (supplementary figure 3d). Fourth, a gene's  
298 repression in salt can be reduced (supplementary figure 3e). Finally, a gene that was  
299 not regulated in the ancestral strain can become regulated in response to salt  
300 (supplementary figure 3f and g). Figure 5a plots the extent to which genes changed in  
301 their regulation against their  $Z_r$  scores. Some of the genes with significant changes in  
302 the regulation are labeled. Figure 6a and b show the five genes whose induction or  
303 repression changed most significantly (based on  $Z_r$ ). We will discuss some of them  
304 further below.

305 There are only four genes in total which showed *increased induction* in the  
306 evolved lines. Among these four genes, two of them (*PUT4* and *PCL5*) also showed  
307 decrease in basal expression in the evolved lines. Four further genes showed *new*  
308 *induction* in the evolved lines. One of them, *CUP1-1* (Figure 5a), has previously been



observed to be up-regulated in response to osmotic stress (Yale & Bohnert, 2001). Five genes showed *reduced induction* in our experiment. One of the genes, *HSP30*, encodes a stress-responsive protein that negatively regulates the H(+)-ATPase Pma1p. It is induced by heat shock, treatment with organic acid and ethanol, and glucose starvation (Panaretou & Piper, 1992; Piper *et al.*, 1994; Piper *et al.*, 1997).

In contrast to the few genes which showed increased induction, a total of 37 genes showed *increased repression*. Out of these 37 genes, one gene (*HLR1*) is directly involved in the stress response. This gene encodes a protein involved in regulation of cell wall composition, as well as in the osmotic stress response (Alonso-Monge *et al.*, 2001). 12 genes showed *new repression* in our experiment. Among these 12 genes, the genes *GCV2* and *GCV1* were previously found to be up-regulated in osmotic stress (Yale & Bohnert, 2001). We did not find any gene with *reduced repression*.

We next classified genes whose regulation changed in any of these ways according to their functions, using the CYGD classification of yeast genes (Güldener *et al.*, 2005) (Figure 7), and compared their distribution among functional categories with the proportions of the yeast genome in each category. The distribution of the number of genes among different classes is non-random for genes whose repression changes (49 genes in total; Chi-square test,  $p\text{-value} < 0.001$ ,  $df=17$ ), whereas the distribution is not significantly different from random for the genes whose induction changes (13 genes;  $p\text{-value} > 0.1$ ,  $df=17$ ). We then asked whether genes whose regulation changes occur preferentially in specific functional categories. Interestingly, for genes with change in induction level, such genes are (marginally) enriched only in the functional class of cell rescue, defense and virulence ( $p=0.0444$ ). Genes whose repression changes are enriched in the functional classes transcription ( $p=0.0006$ ), protein synthesis ( $p=0.0008$ ), and proteins with binding or catalytic function ( $p=0.0003$ ). The classes protein fate (folding, modification, destination;  $p=0.0232$ ), cellular transport ( $p=0.0048$ ), cell cycle/DNA processing ( $p=0.0134$ ) and cell rescue, defense and virulence ( $p=0.0230$ ) contain fewer genes whose repression changes than expected by chance alone (all  $p\text{-values}$  based on an exact binomial test).

#### **Genes with basal level expression change**

Next we turn to genes that show a change in basal expression. There are 82 such genes. Thirty eight of them increased their basal expression, whereas 44 genes reduced their basal expression. Figure 5b plots the extent of change in basal expression ( $\log_2$ -transformed) of all the genes against their corresponding  $Z_b$  values. Some of the genes with significant changes in the basal expression level are labeled. Figure 6c and d lists the top five genes (based on  $Z_b$ ) with increased and decreased basal expression, respectively. The distribution of genes among different functional classes (Figure 7) is significantly different from what would be expected by chance alone for the genes with increase in basal expression ( $p < 0.05$ ,  $df=17$ , Chi-square test), and also for the genes with a decrease in basal expression ( $p < 0.001$ ,  $df=17$ ). For genes with an increase in basal expression, the class 'cell rescue and defense' (stress response) shows significantly more genes than expected ( $p=0.0059$ , exact binomial test), and the class

353 protein fate ( $p=0.0409$ ) shows significantly fewer genes than expected by chance alone.  
354 For genes with a decrease in basal expression, the class 'cell type differentiation' (cell  
355 wall, sporulation, spore wall etc.) ( $p=0.0165$ ) shows more genes than expected, and the  
356 classes 'transcription' ( $p=0.0125$ ) as well as 'protein fate' (folding, modification,  
357 destination) ( $p=0.0014$ ) contain fewer genes than expected.

358 Two well-established stress response genes showed an increase in basal  
359 expression in the evolved lines. One of them is *CTT1*, whose basal expression  
360 increased by ~1.6 fold compared to the ancestral strain. This gene was found to be  
361 induced in four previous genome-wide studies of *physiological* stress adaptation in  
362 yeast (Gasch *et al.*, 2000; Rep *et al.*, 2000; Causton *et al.*, 2001; Yale & Bonhert, 2001).  
363 The second known stress response genes is the transcription factor *MSN4*, whose  
364 expression was increased by approximately 1.5 fold. A more detailed analysis of  
365 individual genes can be found in the supplementary material (supplementary text).

366

#### 367 **Whole Genome (Re)Sequencing Analysis**

368 Some laboratory evolution studies have identified few beneficial mutations of  
369 large fitness effects in evolving populations (Blanc & Adams, 2003; Velicer *et al.*, 2006;  
370 Stanek *et al.*, 2009). To find out if there are any such mutations, we subjected the  
371 ancestral strain and population samples from two of our evolved lines (S1 and S2) to  
372 deep sequencing at about ~10X coverage and the third line (S3) to ~50X coverage,  
373 using the Roche 454 Genome Sequencer. In our analysis of this sequence data, we  
374 aimed to identify only changes that may have swept to high frequency or fixation and  
375 10X coverage is sufficient for that purpose. We chose candidate SNPs based on the  
376 criteria for the changed (derived) nucleotide described in supplementary methods and  
377 our approach is deliberately conservative. We identified 56 candidate SNPs and 7  
378 deletions in total from all the three evolved (S) lines.

379 We sequenced all 56 candidate SNPs and 7 candidate deletions, but only  
380 one of them turned out to be a true change. This change was a SNP unique to line S2,  
381 and occurred at a frequency exceeding 50 percent, based both on next generation  
382 sequencing data and PCR sequencing data. This SNP is a non-synonymous G to A  
383 mutation (amino acid change: G230D) in the *MOT2* (YER068W) gene. The gene *MOT2*  
384 is a subunit of the CCR4-NOT complex which has roles in transcription regulation,  
385 mRNA degradation, and post-transcriptional modifications (Cade & Errede, 1994; Liu *et al.*,  
386 1998; Badarinarayana *et al.*, 2000; Denis *et al.*, 2001; Panasenko *et al.*, 2006;  
387 Mersman *et al.*, 2009), with no known role in salt tolerance. To test whether the *MOT2*  
388 SNP alone provided any fitness benefit to yeast in salt, we replaced the wild-type  
389 variant of *MOT2* in the genome of the ancestral strain with the mutant (see Methods).  
390 Three replicate competition assays using FACS (see Methods) showed a relative  
391 population fitness  $w=1.043\pm0.008$  for the mutant *MOT2* allele (Figure 8a) in medium  
392 with NaCl, compared to  $w=0.999\pm0.014$  for the wildtype *MOT2* allele. However, the  
393 fitness increase of the mutant *MOT2* could only explain 25.3 percent of the total amount  
394 of fitness increase in line S2, whose relative fitness was  $w=1.17$ , indicating that there  
395 are other factors that contribute to the fitness increase of the line S2. In addition, we  
396 note that the *MOT2* mutation also conferred a substantial fitness increase in the

397 absence of NaCl (Figure 8b), indicating that this mutation is not specifically adaptive to  
398 salt stress.

399

#### 400 **Duplication and PFGE**

401 We next turned to copy number variations as sources of evolutionary  
402 adaptation. Since large-scale chromosomal rearrangements and aneuploidies are  
403 frequent in yeast laboratory evolution (Adams *et al.*, 1992; Dunham *et al.*, 2002; Rancati  
404 *et al.*, 2008), and are also observed in yeast gene knock-out strains (Hughes *et al.*,  
405 2000), we first wanted to know whether such rearrangements were numerous in our  
406 evolved lines. To this end, we first performed pulsed field gel electrophoresis (PFGE)  
407 analysis of whole chromosomes for two clones of the ancestral strain, as well as for two  
408 clones from each of the evolved lines. This experiment revealed no changes in any of  
409 the lines, except for one additional band (at ~500kb) that occurred exclusively in line S3  
410 (Figure 9a). PFGE for chromosomes digested with the *NotI* restriction enzyme also  
411 revealed a single novel band (~180kb), which was again unique to line S3 (Figure 9b).  
412 Taken together, these observations suggest that high frequency copy number changes  
413 on a chromosomal scale are not rampant in our evolved lines, and the same  
414 arrangements do not occur across lines. All other distinguishable fragments in the gel,  
415 apart from the novel band of size ~180kb in line S3, were as predicted for a haploid  
416 yeast genome from a computational restriction digest of the genome with *NotI*.  
417 Additionally, chromosome-wide read coverage data from next generation sequencing  
418 (see below) supports the notion that there are no copy number changes involving large  
419 chromosome fragments (supplementary figure 4). However, a change in ploidy in the  
420 evolved lines that would affect entire chromosomes is not detectable by any of these  
421 two methods.

422

#### 423 **Gene Amplifications and Deletions from Next Generation Sequencing data**

424 We next turned to smaller scale copy number changes that we could detect  
425 in our next generation sequencing data. To detect gene duplication and deletions, a  
426 segmentation algorithm (see supplementary methods) was used along with analysis of  
427 DNA breakpoints from the reads (see supplementary Methods). The results of both  
428 analyses are described in detail in the supplementary text.

429 We did not observe any large amplification or deletion in the genomes of any  
430 of the lines (and specifically in line S3) which could explain the appearance of novel  
431 bands in the pulsed field electrophoresis data from Figure 9. This observation, together  
432 with read coverage data of individual chromosomes (Supplementary figure 4) suggests  
433 that these novel bands were results of a non-duplicative translocation event in the line  
434 S3. However, for a haploid genome, a translocation event would displace one of the  
435 original bands in the PFGE, which we did not observe. Thus, such a translocation event  
436 is only consistent with the PFGE data if the affected chromosome occurs in more than  
437 one copy. In other words, it would be consistent with a ploidy change in line S3.

438

#### 439 **Ploidy of the Evolved Yeast Lines**

440 Changes in ploidy of haploid yeast strains under salt stress have been

441 observed before (Gerstein *et al.*, 2006). Neither PFGE nor chromosome wide read  
442 coverage data would be able to detect such changes. To estimate the ploidy level of  
443 evolved lines in comparison to the ancestral strain, we grew ancestral strain and  
444 population samples of the evolved lines for 24 hours at 30°C and estimated the cell  
445 densities. We then isolated genomic DNA from a defined number of cells and quantified  
446 the DNA amount, which allowed us to calculate the DNA content per yeast cell for all  
447 the lines. We found that the DNA content per cell of lines S1 and S2 had increased by  
448 78.7 percent and 85.4 percent, respectively, from that of the ancestral strain. For the  
449 line S3, the DNA content per cell was more than double that of the ancestral strain  
450 (119.91 percent increase from the ancestral strain)(Figure 10). These observations  
451 suggest that all the evolved lines have become massively aneuploid.

452

#### 453 **A consistent cell size increase during our experiment**

454 Increases in ploidy are often associated with cell size increases. Our lines are no  
455 exception, as exemplified by the histogram of Figure 11a, which shows the distribution  
456 of cell diameters of the ancestral strain and the evolved line S2. This increase is also  
457 microscopically visible (Figure 11b). The mean cell diameter increased from 2.35µm  
458 ( $\pm 0.70\mu\text{m}$ ) to 6.73µm ( $\pm 2.34\mu\text{m}$ ). Along with a significant increase in the mean cell  
459 diameter (t-test, p-value  $< 2.2 \times 10^{-16}$ ), the coefficient of variation ( $C_v$ ), as defined by the  
460 ratio of standard deviation and mean, also increased significantly (t-test for distributions  
461 of  $C_v$  for samples drawn from two distributions, p-value  $< 10^{-6}$ ). In other words, not only  
462 did cells become larger during laboratory evolution, they also became more variable in  
463 size. Cell size changes in laboratory evolution experiments are not unprecedented.  
464 They have been observed in evolving *E. coli* populations (Lenski & Travisano, 1994;  
465 Lenski & Mongold, 2000; Philippe *et al.*, 2009), as well as in *Staphylococcus aureus*  
466 populations in NaCl medium (Vijaranakul *et al.*, 1995). However, effect of cell size on  
467 the survival and fitness are poorly understood.

468

#### 469 **Discussion**

470 To investigate evolutionary adaptation to long-term osmotic stress, we  
471 evolved yeast cells in the laboratory for 300 generations in high salt (NaCl) medium. We  
472 observed that salt reduces the growth rate of our ancestral strain by 11 percent.  
473 Consequently, the final cell density after 24 hours of growth is approximately 3 times  
474 lower compared to cells grown in the same medium but without salt. Our three replicate  
475 evolved lines grow approximately 8 to 12 percent faster than the ancestral strain in high  
476 salt medium, and part of this increase reflects adaptation specific to salt.

477 We analyzed the gene expression levels in all the evolved lines as well as in  
478 the ancestral strain. Although there were many shared genes between the evolved lines  
479 and the ancestral strain that respond to salt in similar manner, we also observed  
480 multiple differentially expressed genes in the evolved lines compared to the ancestral  
481 strain. These differentially expressed genes can be divided into two categories. The first  
482 category comprises genes with changes in their basal expression level in the evolved  
483 lines compared to the ancestor (i.e., even in the absence of salt). The second category  
484 comprises genes regulated differently in response to salt in the evolved lines.

485 Multiple genes whose expression shows an *evolutionary* response in our  
486 experiments were also affected by hyperosmotic stress or salt stress in previous studies  
487 of the *physiological* stress response. Changed expression of many genes in the  
488 *physiological* stress response in general, and in the hyperosmotic stress response in  
489 particular is transient (Gasch *et al.*, 2000; Causton *et al.*, 2001), and may depend on the  
490 time and conditions in which it is measured. It is thus to be expected that different  
491 studies show limited comparability with respect to the identity of these genes.

492 Some of the genes whose expression evolved are known to be associated  
493 with the stress response. For example, two genes, *CTT1* and *GAC1*, showed an  
494 increase in basal expression in the evolved lines compared with the ancestral strain.  
495 Both these genes were also up-regulated in response to salt in our ancestral strain.  
496 *CTT1* encodes for a cytosolic catalase and protects the cell from oxidative damage by  
497 reactive oxygen species (Jamieson, 1998; Lushchak & Gospodaryov, 2005; Herrero *et al.*, 2008). This gene has been shown to be up-regulated in four previous studies of  
498 *physiological* stress response and salt stress response (Gasch *et al.*, 2000; Rep *et al.*,  
500 2000; Causton *et al.*, 2001; Yale & Bohnert, 2001). The gene *GAC1* is associated with  
501 glycogen synthesis (Wu *et al.*, 2001) and is also induced during osmo-adaptation  
502 (Posas *et al.*, 2000). Both these genes contain stress responsive elements (STRE) in  
503 their promoters (Schuller *et al.*, 1994; Moskvina *et al.*, 1998) and are known to be  
504 activated under various stress conditions. A third gene, *CUP1-1*, showed increased  
505 induction in the evolved lines and also was up-regulated in response to NaCl in our  
506 ancestral strain. This gene was also shown to be up-regulated in one previous study of  
507 *physiological* stress response (Yale & Bohnert, 2001). Yet another gene, *HLR1*, showed  
508 increased repression in the evolved lines, and was down-regulated in the ancestral  
509 strain in NaCl medium. This gene encodes a protein involved in maintaining cell wall  
510 composition and is also involved in response to osmotic stress (Alonso-Monge *et al.*,  
511 2001).

512 There are three main *physiological* components of the salt stress response.  
513 They affect adaptation time to salt, growth rate in salt, and efficiency of growth in salt  
514 (Warringer *et al.*, 2003). In our experiments, *evolutionary* adaptation to salt must affect  
515 one or more of these three component, because the viability itself of yeast cells does  
516 not change between the ancestral strain and the evolved lines. There are various ways  
517 in which these components could change during evolution. Firstly, some of the genes  
518 whose basal expression changes might be directly involved in the initial adaptation to  
519 salt. Increasing or decreasing the basal expression of those genes to the level needed  
520 for salt adaptation, might enable the cells respond to salt much more quickly as  
521 opposed to changing their expression state. For example, some ion transporter genes  
522 (e.g. *FRE5*) have an increased basal expression in the evolved lines in our experiment.  
523 Higher levels of these transporter proteins at the initial stages of the salt stress  
524 response would help cells achieve ion homeostasis much more quickly. Secondly, some  
525 of the differentially expressed genes might not be directly related to the salt stress  
526 response but encode transcription factors that controls salt stress response genes. For  
527 example, the transcription factor *MSN4*, that controls many stress response genes, has  
528 increased basal expression levels in our evolved yeast lines. Higher levels of this

transcription factor during the initial adaptation phase could ensure faster induction of the genes required for salt stress response. Thirdly, there are genes that show evolutionary change in regulation in the evolved lines. Higher level of induction or repression of these genes might increase the growth rate and/or efficiency of growth in salt medium.

We found 16 genes that gained regulation (new induction and new repression) in the evolution experiment, suggesting that new ways of salt stress adaptation could also arise during evolution. However, we also observed loss of regulation, at least to some extent, for five genes (genes with reduced induction and reduced repression) in the evolved lines. Because we performed our evolution experiment under constant environmental stress, losing regulation might be advantageous for evolutionary adaptation to salt stress. If the affected genes are involved in adaptation to multiple stresses, losing regulation could actually be beneficial both under constant or fluctuating environmental stressors. However, if the affected genes are specific for that particular stress, a loss in regulation may also have a cost, because cells would become physiologically less flexible under fluctuating environmental stresses. Among the five genes with loss of regulation, one gene, *HSP30*, can be induced by several stresses (Piper *et al.*, 1997). However, the mRNA level of this gene was shown to be diminished in NaCl medium by Rep *et al* (Rep *et al.*, 2000).

The hyperosmotic and salt stress responses are complex and influenced by many genes. It is an open question whether a small number of genetic changes with large effects could dramatically increase salt stress resistance, and thus explain the fitness increase in our evolved lines. To find out, we also sequenced the ancestral strain and population samples of the evolved lines using deep sequencing with genome coverages between 10- and 50-fold to identify whether any high frequency polymorphisms arose in the evolved lines during our evolution experiment. For two reasons, we chose deep population sequencing over clone sequencing. First, population sequencing gives a comprehensive view of polymorphisms in a population. Second, it can also permit estimation of polymorphism frequencies, although any such estimate would be more qualitative than quantitative at our sequence coverage. At the very least, population sequencing can detect high frequency polymorphisms. Such polymorphisms are of the greatest interest to us, because they would correspond to adaptive mutations with strong fitness effects that rose to high frequencies. We note that the population sizes in our experiment are so large that no neutral polymorphisms would be expected to rise to high frequencies during the experiment's short duration except through hitchhiking with an advantageous mutation (effective population size in our experiment is  $>2 \times 10^6$ ). (Clone sequencing can detect low frequency polymorphisms, but for a heterogeneous population, this method requires sequencing of many clones, and is thus currently prohibitively costly.)

Only one of our evolved lines (S2) contained a high frequency single nucleotide polymorphism (SNP). This SNP occurs in more than 50 percent of the population, and causes an amino acid substitution in the protein encoded by the *MOT2* gene. When introduced into the ancestral strain, this mutation causes a fitness increase

573 that explains 25.3 percent of the increase in fitness  $w$  in the evolved strain. However,  
574 the mutation causes a fitness increase also in the absence of salt, which means that its  
575 effects do not reflect a specific adaptation to salt.

576 **Multiple genetic changes of low population frequency could be present in our**  
577 **evolved lines, and these changes might explain a part of the fitness increase we see.**

578 Such low frequency SNPs might be present for several reasons. First, individual  
579 mutations may confer only modest fitness benefits and thus increase in frequency  
580 slowly. Second, some mutations with very strong fitness effects may have occurred, but  
581 late in the experiment, and thus might simply not have had enough time to rise to high  
582 frequency. Third, the fitness effect of alleles might be determined by epistatic  
583 interactions with other mutations elsewhere in the genome (Elena & Lenski, 1997; Elena  
584 & Lenski, 2001; Morgan & Feldman, 2001). Eventually genotypes with such alleles  
585 might rise to high frequency, but at time scales much longer than those of our  
586 experiment, because multiple mutations might have to occur before any one epistatic  
587 combination with strong benefits arises. Finally, and perhaps most importantly, there  
588 could be clonal interference between multiple beneficial polymorphisms in the  
589 population and this might prevent any particular polymorphism from rising to high  
590 frequency in the population (Kao & Sherlock, 2008).

591 Similar to our analysis on SNPs, our analysis on copy number variations,  
592 both based on PFGE and next generation sequencing, did not reveal any large scale  
593 copy number changes shared across strains. However, we observed ploidy increases in  
594 all the evolved lines, and this could be the reason behind increase in cell size of these  
595 lines.

596 Evidence from plants suggests that an increase in ploidy and the resultant  
597 increase in cell size can be advantageous under salt-stressed conditions. For example,  
598 within a species, plants with higher ploidy can cope with salt stress better than plants  
599 with lower ploidy (Saleh *et al.*, 2008). Polyploid plants have higher water content and  
600 lower osmotic pressure than diploid plant (Noggle, 1946). Their increased water content  
601 is due to a decrease in the surface-to-volume ratio of cells (Stebbins, 1950). In addition,  
602 any water content decrease due to high salinity is smaller in polyploid plants than that in  
603 diploid plants (Tal & Gardi, 1976). Gerstein *et al.* observed that initially haploid yeast  
604 lines increased genome size significantly faster under salt stress than in unstressed  
605 condition, suggesting an advantage for higher ploidy or increased cell size also in yeast  
606 (Gerstein *et al.*, 2006).

607 Changes in ploidy could help yeast cells adapt faster physiologically, for  
608 example by affecting expression of genes important for salt stress tolerance. Could all  
609 the changes in gene expression that we observed in our experiment be caused solely  
610 by ploidy changes or by the concomitant increase in cell size? To find out, we compared  
611 the set of genes that changed expression in our evolved lines with the set of genes that  
612 are known to change expression after cell size increases (Wu *et al.*, 2010) and ploidy  
613 changes (Galitski *et al.*, 1999). Remarkably, only three genes that evolve changed  
614 expression in our experiment (out of 144 differentially expressed genes in our dataset)  
615 are among the genes affected by cell size increase or ploidy changes. Two of these  
616 genes showed increased basal expression in our experiment. One of them, YIL169C,

617 was also observed to be induced with increase in cell size (Wu *et al.*, 2010). The other  
618 gene, *COS8*, was observed to change expression with a change in ploidy level (Galitski  
619 *et al.*, 1999). The third gene, *YLR042C*, showed a basal decrease in expression in our  
620 evolved lines. It was also observed to be repressed in cells with increased size (Wu *et*  
621 *al.*, 2010).

622 Overall, the vast majority of changes that we observe in gene expression  
623 cannot be caused by changed ploidy or cell size. Because our whole genome  
624 sequencing analysis revealed no small genetic changes of strong phenotypic effects,  
625 the gene expression changes we observe could be caused by multiple genetic changes  
626 of modest individual effects. Some of these changes might affect a gene's expression in  
627 *cis*, others in *trans*, for example through changes in the amino acid sequence of  
628 transcriptional regulators (Wittkopp *et al.*, 2004; Emerson & Li, 2010). In addition, some  
629 expression changes may be caused by epigenetic change (e.g., changes in DNA  
630 methylation) or alterations in cellular memory (Turner, 2002; Ringrose & Paro, 2004;  
631 Zhang *et al.*, 2005; Zacharioudakis *et al.*, 2007).

632 We found several genes with modest changes in expression level in our  
633 evolved lines, and no gene with very strong expression change. Our observations stand  
634 in contrast to the only laboratory evolution study we know of that investigated  
635 evolutionary adaptation to a stressor. The study asked how *E. coli* cells adapt to heat  
636 stress. It observed very high level of evolutionary expression change in two proteins,  
637 GroEL and GroES, that are known heat shock genes (Rudolph *et al.*, 2010). The reason  
638 for this difference to our work is probably that adaptation to salt stress requires several  
639 molecular functions simultaneously, namely ionic detoxification to achieve ionic  
640 equilibrium, maintaining cellular water activity, and osmolyte synthesis. Each of the  
641 evolutionarily altered genes in our experiment may perform or control some of these  
642 functions and thus, may have only a modest individual fitness contribution when  
643 differentially expressed. In addition, gene interactions (Phillips 2008; He *et al.*, 2010)  
644 could also contribute towards fitness increase. This scenario resembles recent  
645 observations in genome wide association studies (GWAS) for complex human traits and  
646 diseases, where researchers have observed many genetic variants with modest effects  
647 on a phenotype (Frayling, 2007; Barrett *et al.*, 2008; Cooper *et al.*, 2008; Visscher,  
648 2008; Zeggini *et al.*, 2008; Hindorff *et al.*, 2009; Manolio *et al.*, 2009; Visscher &  
649 Montgomery, 2009; Park *et al.*, 2010; Yang *et al.*, 2010).

650 Taken together, adaptation to salt stress is associated with DNA content  
651 increase and gene expression change in all the three lines, one high frequency mutation  
652 in one of the evolved lines, and one chromosomal rearrangement in another. The  
653 increase in DNA content in all the lines is massive and previous studies suggest that the  
654 change in ploidy might be beneficial for the lines in salt. In contrast, there are 144 genes  
655 in our study that show modest change in their expression levels. This observation  
656 suggests that tinkering with expression levels of several genes might be a useful  
657 mechanism to respond to a complex, polygenic trait rather than altering expression of  
658 few genes to very high levels. Such tinkering might also reflect cellular constraints on  
659 evolution. For example, changing expression of a gene to very high levels or  
660 suppressing a gene completely might increase fitness of the yeast cells in salt medium.



661 However, changing expression of a gene to very high levels or suppressing a gene  
662 completely might affect the capability of the cells to grow in other media. Since we do  
663 not find high frequency genetic changes that can explain expression changes of the  
664 genes, it is possible that these are caused by multiple genetic changes of low frequency  
665 (that might vary from one subpopulation to other) or by epigenetic changes. These  
666 results also suggest that cells in our populations perhaps employ several strategies in  
667 combination for adaptation to salt stress rather than following a single path to  
668 adaptation. It is possible that such genetic heterogeneity might render the populations to  
669 be more robust in the long term. If some of the genetic factors contributing to the salt  
670 stress adaptation are lost, the fitness of the population would not fall drastically. Also,  
671 such diverse populations might be less vulnerable against other environmental  
672 fluctuations which might co-occur with saline stress in natural environment. Although  
673 individual contributions of each of these genetic mechanisms to adaptation remain  
674 unclear, our study provides a stepping stone for asking further questions on  
675 *evolutionary* adaptation of yeast to salt stress.

676

## 677 **Methods**

### 678 **Strains and media.**

679 All laboratory evolution experiments started from the same clone of haploid yeast strain  
680 BY4741, which is referred to as the *ancestral* strain. The 3 replicate yeast lines evolved  
681 in NaCl are referred to as lines S1, S2 and S3. The growth rates of the evolved lines as  
682 well as the ancestral strain were measured relative to a BY4741 strain in which the  
683 *CWP2* gene was GFP-tagged (termed as the *reference* strain). For serial transfers, cells  
684 were cultured in YP and 2% galactose (YPG) and YPG supplemented with 0.5 M NaCl  
685 (YPGN).

686

### 687 **Serial transfer**

688 Six parallel serial transfer experiments were started from one single clone of the  
689 ancestral strain. In each parallel experiment, 50 ml yeast culture was grown for at 30°C.  
690 Every 24 hours, 50µl of grown culture was transferred into fresh culture medium; 30  
691 such transfer cycles were carried out for a total of approximately 300 generations (Each  
692 transfer cycle involved approximately  $\log_2 1000 \approx 10$  cell generations). In three of the  
693 parallel experiments, the culture medium was YPG, whereas in the other three parallel  
694 experiments, the medium was YPGN.

695

### 696 **Viability assays**

697 To estimate cell viabilities, cultures of growing yeast cells were sampled after 16 hours  
698 (during exponential growth phase), as well as after 24 hours (during stationary phase),  
699 diluted and plated. The plates were incubated at 30°C for 5 days and the number of  
700 colonies were counted. The relative viabilities of both the ancestral strain and the  
701 evolved lines were estimated from the ratio of colonies forming on agar plates  
702 containing YPG+ 0.5M NaCl to that of plates containing only YPG. All the  
703 measurements were carried out in three biological replicates.

704

## 705 **Competition assays**

706 To compare growth rates of the evolved lines with that of the ancestral strain,  
707 competition assays were carried out in triplicates using fluorescence activated cell  
708 sorting (FACS). Cells from frozen glycerol stocks were grown overnight in 4ml YPD  
709 medium (30°C, 220rpm) until late logarithmic phase ( $<1.5 \times 10^8$  cells/ml). For the  
710 competition assay, approximately equal cell numbers of the reference strain and of the  
711 competing strain were mixed and grown for 24 hours at 30°C. The relative cell numbers  
712 at the beginning and at the end of the competition experiment were determined using  
713 FACS (supplementary methods) and growth rate differences were estimated as  
714 described in supplementary methods. **For calculation of fitness, the relative cell**  
715 **numbers at the beginning of the assay were taken into consideration.**

716

## 717 **Whole genome transcriptome analysis**

718 The mRNA expression levels in the ancestral strain and in the evolved lines were  
719 analyzed using a GeneChip Yeast Genome 2.0 Array (Affymetrix). Equal number of  
720 cells from the ancestral strain and the evolved lines were grown in YPG medium for 16  
721 hours. The cells were then either induced with 0.5M NaCl or grown uninduced for 20  
722 further minutes. The microarray analysis was carried out for 2 replicates each for the  
723 ancestral strain in YPG and YPGN (4 in total) and for 4 replicate population samples (1  
724 for S1, 2 for S2 and 1 for S3) each in YPG and YPGN for the S lines (8 in total).  
725 “Shared” genes, genes that respond to NaCl in a similar manner between the ancestral  
726 strain and the evolved lines, with significant up-regulation or down-regulation were  
727 identified using a t-test at  $p=0.05$ , False Discovery Rate (FDR)  $<10\%$  and  $|\log_2(\text{fold}$   
728  $\text{change})| \geq 1$ .

729 Genes with changes in basal expression or change in regulation were  
730 identified based on two Z scores  $Z_r$  and  $Z_b$  (see supplementary methods). Genes whose  
731 absolute Z-scores exceeded a value of 1.5 were considered to be differentially  
732 expressed. The differentially expressed genes were then grouped into different classes  
733 using the CYGD functional classification for yeast genes (Guldener *et al.*, 2005).

734

## 735 **Whole Genome Sequencing and SNP identification**

736 The ancestral strain, as well as population samples of the NaCl evolved lines at  
737 generation 300 were sequenced at approximately 10X coverage using next generation  
738 pyrosequencing (Margulies *et al.*, 2005) (genome sequencer FLX LR system, Roche).  
739 The line S3 was further sequenced to a total of ~50X coverage. Candidate SNPs and  
740 indels were identified using blastn (Altschul *et al.*, 1990) followed by MUSCLE (Edgar,  
741 2004) based on the criteria described in supplementary methods. PCR sequencing was  
742 done to confirm candidate SNPs and indels.

743

## 744 **Pulsed-field gel electrophoresis (PFGE)**

745 To identify large scale chromosomal rearrangements, two clones from each of the  
746 evolved lines (S1, S2 and S3) and the ancestral strain were analyzed by PFGE.  
747 Agarose plugs were prepared with the CHEF Yeast Genomic DNA Plug Kit (Bio-Rad)  
748 according to the manufacturer's protocol. For restriction enzyme digestion, agarose

749 plugs were incubated with NotI restriction enzyme for 16hrs at 37°C mixture in an  
750 appropriate restriction buffer. The digested or undigested plugs were then loaded into  
751 the wells of 1% agarose gels. Electrophoresis was carried out in a CHEF-DRIII system  
752 (Bio-Rad). The gels were stained using ethidium bromide (EtBr) and photographed.

753

#### 754 **Gene Amplification and Deletions from Next Generation Sequencing data**

755 In addition to identifying very short indels within individual reads, larger gene  
756 duplications and deletions in the evolved lines, compared to the ancestral strain, were  
757 identified using read coverage information from the sequencing data. Furthermore, DNA  
758 breakpoint analysis was performed to confirm those candidate copy number changes.

759

#### 760 **Ploidy determination of the evolved lines**

761 To estimate the ploidy level of the evolved lines in comparison to the  
762 ancestral strain, the ancestral strain and population samples of the evolved lines were  
763 grown for 24 hours at 30°C. Genomic DNA was isolated from a defined number of cells  
764 following the protocol described in (Sambrook & Russell, 2001, Chapter 6).

765

#### 766 **Cell size measurement**

767 Population samples of ancestral and evolved cells were grown in YPG medium up to  
768 mid-log phase. The cell density was adjusted to be the same for all the samples  
769 (~1x10<sup>7</sup> cells/ml). 20ul of cells were then loaded onto disposable counting chambers and  
770 the cell sizes, were measured using a Cellometer M10 (Nexcelom Biosciences),  
771 following the manufacturer's protocol.

772

#### 773 **Acknowledgements**

774 We would like to acknowledge the help of the Functional Genomics Center Zurich for  
775 carrying out next generation sequencing and microarray experiments. AW  
776 acknowledges support through SNF grants 315200-116814, 315200-119697, and  
777 315230-129708, as well as through the YeastX project of SystemsX.ch, and the  
778 University Research Priority Program in systems biology at the University of Zurich.

#### 779 **References**

- 780 Adams, J. & Oeller, P.W. 1986. Structure of evolving populations of *Saccharomyces*  
781 *cerevisiae*: Adaptive changes are frequently associated with sequence alterations  
782 involving mobile elements belonging to the Ty family. *Proc. Natl. Acad. Sci. USA* **83**:  
783 7124-7127.
- 784
- 785 Adams, J., Puskas-Rozsa, S., Simlar, J. & Wilke, C.M. 1992. Adaptation and major  
786 chromosomal changes in populations of *Saccharomyces cerevisiae*. *Curr. Genet.* **22**:  
787 13-19.
- 788
- 789 Alonso-Monge, R., Real, E., Wojda, I., Bebelman, J.P., Mager, W.H. & Siderius, M.  
790 2001. Hyperosmotic stress response and regulation of cell wall integrity in  
791 *Saccharomyces cerevisiae* share common functional aspects. *Mol. Microbiol.* **41**: 717-

792 730.  
793  
794 Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. 1990. Basic local  
795 alignment search tool. *J. Mol. Biol.* **215**: 403-410.  
796  
797 Apse, M.P., Aharon, G.S., Snedden, W.A. & Blumwald, E. 1999. Salt Tolerance  
798 Conferred by Overexpression of a Vacuolar Na<sup>+</sup>/H<sup>+</sup> Antiport in *Arabidopsis*. *Science*  
799 **285**: 1256-1258.  
800  
801 Attfield, P.V. 1997. Stress Tolerance: The key to effective strains of industrial baker's  
802 yeast. *Nat. Biotech.* **15**: 1351-1357.  
803  
804 Badarinarayana, V., Chiang, Y.C. & Denis, C.L. 2000. Functional interaction of CCR4-  
805 NOT proteins with TATAA-binding protein (TBP) and its associated factors in yeast.  
806 *Genetics* **155**: 1045-1054.  
807  
808 Barrett, J.C., Hansoul, S., Nicolae, D.L., Cho, J.H., Duerr, R.H., Rioux, J.D. *et al.* 2008.  
809 Genome-wide association defines more than 30 distinct susceptibility loci for Crohn's  
810 disease. *Nat. Genet.* **40**: 955-962.  
811  
812 Betz, C., Zajonc, D., Moll, M. & Schweizer, E. 2002. ISC1-encoded inositol  
813 phosphosphingolipid phospholipase C is involved in Na<sup>+</sup>/Li<sup>+</sup> halotolerance of  
814 *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **269**: 4033-4039.  
815  
816 Blanc, V.M. & Adams, J. 2003. Evolution in *Saccharomyces cerevisiae*: Identification of  
817 Mutations Increasing Fitness in Laboratory Populations. *Genetics* **165**: 975-983.  
818  
819 Bressan, R.A., Hasegawa, P.M. & Pardo J.M. 1998. Plants use calcium to resolve salt  
820 stress. *Trends Plant Sci.* **3**: 411-412.  
821  
822 Brewster, J.L., de Valoir, T., Dwyer, N.D., Winter, E. & Gustin, M.C. 1993. An  
823 Osmosensing Signal Transduction Pathway in Yeast. *Science* **259**: 1760-1763.  
824  
825 Cade, R.M. & Errede, B. 1994. MOT2 encodes a negative regulator of gene expression  
826 that affects basal expression of pheromone-responsive genes in *Saccharomyces*  
827 *cerevisiae*. *Mol. Cell. Biol.* **14**: 3139-3149.  
828  
829 Causton, H.C., Ren, B., Koh, S.S., Harbison, C.T., Kanin, E., Jennings, E.G. *et al.* 2001.  
830 Remodeling of Yeast Genome Expression in Response to Environmental Changes. *Mol.*  
831 *Biol. Cell* **2**: 323-337.  
832  
833 Cooper, J.A. 1994. MAP KINASE PATHWAYS: Straight and narrow or tortuous and  
834 intersecting? *Curr. Biol.* **4**: 1118-1121.  
835

836 Cooper, J.D., Smyth, D.J., Smiles, A.M., Plagnol, V., Walker, N.M., Allen, J.E. *et al.*  
837 2008. Meta-analysis of genome-wide association study data identifies additional type 1  
838 diabetes risk loci. *Nat. Genet.* **40**: 1399-1401.

839

840 Denis, C.L., Chiang, Y.C., Cui, Y. & Chen, J. 2001. Genetic evidence supports a role for  
841 the yeast CCR4-NOT complex in transcriptional elongation. *Genetics* **158**: 627-634.

842

843 Dihazi, H., Kessler, R. & Eschrich, K. 2004. High Osmolarity Glycerol (HOG) Pathway-  
844 induced Phosphorylation and Activation of 6-Phosphofructo-2-kinase Are Essential for  
845 Glycerol Accumulation and Yeast Cell Proliferation under Hyperosmotic Stress. *J. Biol.*  
846 *Chem.* **279**: 23961-23968.

847

848 Dunham, M.J., Badrane, H., Ferea, T., Adams, J., Brown, P.O., Rosenzweig, F. *et al.*  
849 2002. Characteristic genome rearrangements in experimental evolution of  
850 *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **99**: 16144-16149.

851

852 Edgar, R.C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high  
853 throughput. *Nucleic Acids Res.* **32**:1792-1797.

854

855 Elena, S.F. & Lenski, R.E. 1997. Test of synergistic interactions among deleterious  
856 mutations in bacteria. *Nature* **390**: 395-398.

857

858 Elena, S.F. & Lenski, R.E. 2001. Epistasis between new mutations and genetic  
859 background and a test of genetic canalization. *Evolution* **55**: 1746-1752.

860

861 Emerson, J.J. & Li, W-H. 2010. The genetic basis of evolutionary change in gene  
862 expression levels. *Phil. Trans. R. Soc. B* **365**: 2581-2590.

863

864 Errede, B. & Levin, D.E. 1993. A conserved kinase cascade for MAP kinase activation  
865 in yeast. *Curr. Opin. Cell Biol.* **5**: 254-260.

866

867 Espinosa-Ruiz, A., Belles, J.M., Serrano, R. & Culianez-Macla, V. 1999. *Arabidopsis*  
868 *thaliana* AtHAL3: A flavoprotein related to salt and osmotic tolerance and plant growth.  
869 *Plant J.* **20**: 529-539.

870

871 Ferea, T.L., Botstein, D., Brown, P.O. & Rosenzweig, R.F. 1999. Systematic changes in  
872 gene expression patterns following adaptive evolution in yeast. *Proc. Natl. Acad. Sci.*  
873 *USA.* **96**: 9721-9726.

874

875 Ferrando, A., Kron, S.J., Rios, G., Fink, G.R. & Serrano, R. 1995. Regulation of cation  
876 transport in *Saccharomyces cerevisiae* by the salt tolerance gene HAL3. *Mol. Cell. Biol.*  
877 **15**: 5470-5481.

878

879 Frayling, T.M. 2007. Genome-wide association studies provide new insights into type 2

880 diabetes aetiology. *Nat. Rev. Genet.* **8**: 657-662.

881

882 Galcheva-Gargova, Z., Derijard, B., Wu, I.H. & Davis, R.J. 1994. An osmosensing signal  
883 transduction pathway in mammalian cells. *Science* **265**: 806-808.

884

885 Galitski, T., Saldanha, A.J., Styles, C.A., Lander, E.S. & Fink, G.R. 1999. Ploidy  
886 regulation of gene expression. *Science* **285**: 251-254.

887

888 Ganster, R.W., McCartney, R.R. & Schmidt, M.C. 1998. Identification of a Calcineurin-  
889 Independent Pathway Required for Sodium Ion Stress Response in *Saccharomyces*  
890 *cerevisiae*. *Genetics* **150**: 31-42.

891

892 Garciadeblas, B., Rubio, F., Quintero, F.J., Bañuelos, M.A., Haro, R. & Rodríguez-  
893 Navarro, A. 1993. Differential expression of two genes encoding isoforms of the ATPase  
894 involved in sodium efflux in *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* **236**: 363-368.

895

896 Gasch, A.P., Spellman, P.T., Kao, C.M., Carmel-Harel, O., Eisen, M.B., Storz, G. *et al.*  
897 2000. Genomic Expression Programs in the Response of Yeast Cells to Environmental  
898 Changes. *Mol. Biol. Cell* **11**: 4241-4257.

899

900 Gaxiola, R., de Larrinoa, I.F., Villalba, J.M. & Serrano, R. 1992. A novel and conserved  
901 salt-induced protein is an important determinant of salt tolerance in yeast. *EMBO J.* **11**:  
902 3157-3164.

903

904 Gaxiola, R.A., Rao, R., Sherman, A., Grisafi, P., Alper, S.L. & Fink, G.R. 1999. The  
905 *Arabidopsis thaliana* proton transporters, AtNhx1 and Avp1, can function in cation  
906 detoxification in yeast. *Proc. Natl. Acad. Sci. USA* **96**: 1480-1485.

907

908 Gerstein, A.C., Chun, H.J.E., Grant, A. & Otto, S.P. 2006. Genomic convergence toward  
909 diploidy in *Saccharomyces cerevisiae*. *PLoS Genet.* **2**: e145.

910

911 Gisbert, C., Rus, A.M., Bolarín, M.C., Lopez-Coronado, J.M., Arrillaga, I., Montesinos,  
912 C. *et al.* 2000. The Yeast HAL1 Gene Improves Salt Tolerance of Transgenic Tomato.  
913 *Plant Physiol.* **123**: 393-402.

914

915 Goossens, A., Forment, J. & Serrano, R. 2002. Involvement of Nst1p/YNL091w and  
916 Msl1p, a U2B" splicing factor, in *Saccharomyces cerevisiae* salt tolerance. *Yeast* **19**:  
917 193-202.

918

919 Gorner, W., Durchschlag, E., Martinez-Pastor, M.T., Estruch, F., Ammerer, G. &  
920 Hamilton, B. *et al.* 1998. Nuclear localization of the C2H2 zinc finger protein Msn2p is  
921 regulated by stress and protein kinase A activity. *Genes Dev.* **12**: 586-597.

922

923 Gresham, D., Desai, M.M., Tucker, C.M., Jenq, H.T., Pai, D.A., Ward, A. *et al.* 2008.

924 The Repertoire and Dynamics of Evolutionary Adaptations to Controlled Nutrient-  
 925 Limited Environments in Yeast. *PLoS Genet.* **4**: e1000303.  
 926  
 927 Güldener, U., Münsterkötter, M., Kastenmüller, G., Strack, N., van Helden, J., Lemer, C.  
 928 *et al.* 2005. CYGD: the Comprehensive Yeast Genome Database. *Nucl. Acids Res.* **33**:  
 929 D364-D368.  
 930  
 931 Han, J., Lee, J.D., Bibbs, L. & Ulevitch, R.J. 1994. A MAP kinase targeted by endotoxin  
 932 and hyperosmolarity in mammalian cells. *Science* **265**: 808-811.  
 933  
 934 Haro, R., Garciadeblas, B. & Rodríguez-Navarro, A. 1991. A novel P-type ATPase from  
 935 yeast involved in sodium transport. *FEBS Lett.* **291**: 189-191.  
 936  
 937 Hasegawa, P.M. & Bressan, R.A. 2000. Plant Cellular and Molecular Responses to  
 938 High Salinity. *Annu. Rev. Plant. Physiol. Plant. Mol. Biol.* **51**: 463-499.  
 939  
 940 He, X., Qian, W., Wang, Z., Li, Y. & Zhang, J. 2010. Prevalent positive epistasis in  
 941 *Escherichia coli* and *Saccharomyces cerevisiae* metabolic networks. *Nat. Genet.* **42**:  
 942 272-276.  
 943  
 944 Heath, V.L., Shaw, S.L., Roy, S. & Cyert, M.S. 2004. Hph1p and Hph2p, Novel  
 945 Components of Calcineurin-Mediated Stress Responses in *Saccharomyces cerevisiae*.  
 946 *Eukaryotic Cell* **3**: 695-704  
 947  
 948 Herrero, E., Ros, J., Bellí, G. & Cabisco, E. 2008. Redox control and oxidative stress in  
 949 yeast cells. *Biochim. Biophys. Acta* **1780**: 1217-1235.  
 950  
 951 Hindorff, L.A., Sethupathy, P., Junkins, H.A., Ramos, E.M., Mehta, J.P., Collins, F.S. *et*  
 952 *al.* 2009. Potential etiologic and functional implications of genome-wide association loci  
 953 for human diseases and traits. *Proc. Natl. Acad. Sci. USA* **106**: 9362–9367.  
 954  
 955 Hirata, Y., Andoh, T., Asahara, T. & Kikuchi, A. 2003. Yeast Glycogen Synthase Kinase-  
 956 3 Activates Msn2p-dependent Transcription of Stress Responsive Genes. *Mol. Biol.*  
 957 *Cell.* **14**: 302-312.  
 958  
 959 Hirt, H. 1997. Multiple roles of MAP kinases in plant signal transduction. *Trends Plant*  
 960 *Sci.* **2**: 11-15.  
 961  
 962 Hohmann, S. 2002. Osmotic Stress Signaling and Osmoadaptation in Yeasts. *Micro.*  
 963 *Mol. Biol. Rev.* **66**: 300–372.  
 964  
 965 Hughes, T.R., Roberts, C.J., Dai, H., Jones, A.R., Meyer, M.R., Slade, D. *et al.* 2000.  
 966 Widespread aneuploidy revealed by DNA microarray expression profiling. *Nat. Genet.*  
 967 **25**: 333-337.

968  
 969 Jamieson, D.J. 1998. Oxidative stress responses of the yeast *Saccharomyces*  
 970 *cerevisiae*. *Yeast* **14**: 1511-1527.  
 971  
 972 Jonak, C., Kiegerl, S., Ligterink, W., Barker, P.J., Huskisson, N.S. & Hirt, H. 1996.  
 973 Stress signaling in plants: A mitogen-activated protein kinase pathway is activated by  
 974 cold and drought. *Proc. Natl. Acad. Sci. USA* **93**: 11274-11279.  
 975  
 976 Jonak, C., Ligterink, W. & Hirt, H. 1999. MAP kinases in plant signal transduction. *Cell.*  
 977 *Mol. Life Sci.* **55**: 204-213.  
 978  
 979 Kao, K.C. & Sherlock, G. 2008. Molecular characterization of clonal interference during  
 980 adaptive evolution in asexual populations of *Saccharomyces cerevisiae*. *Nat. Genet.* **40**:  
 981 1499-1504.  
 982  
 983 Kosako, H., Nishida, E. & Gotoh, Y. 1993. cDNA cloning of MAP kinase kinase reveals  
 984 kinase cascade pathways in yeasts to vertebrates. *EMBO J.* **12**: 787-794.  
 985  
 986 Kovtun, Y., Chiu, W-L., Tena, G., Sheen, J. 2000. Functional analysis of oxidative  
 987 stress-activated mitogen-activated protein kinase cascade in plants. *Proc. Natl. Acad.*  
 988 *Sci. USA* **97**: 2940-2945.  
 989  
 990 Kyriakis, J.M. & Avruch, J. 2001. Mammalian Mitogen-Activated Protein Kinase Signal  
 991 Transduction Pathways Activated by Stress and Inflammation. *Physiol. Rev.* **81**: 807-  
 992 869.  
 993  
 994 Lee, J.H., Montagu, M.V. & Verbruggen, N. 1999. A highly conserved kinase is an  
 995 essential component for stress tolerance in yeast and plant cells. *Proc. Natl. Acad. Sci.*  
 996 *USA* **96**: 5873-5877.  
 997  
 998 Lenski, R.E. & Travisano, M. 1994. Dynamics of adaptation and diversification: A  
 999 10,000-generation experiment with bacterial populations. *Proc. Natl. Acad. Sci. USA* **91**:  
 1000 6808-6814.  
 1001  
 1002 Lenski, R.E. & Mongold, J.A. 2000. Cell size, shape, and fitness in evolving populations  
 1003 of bacteria. In: *Scaling in Biology* (J. H. Brown & G. B. West, eds), pp. 221-235, Oxford  
 1004 University Press.  
 1005  
 1006 Liu, H.Y., Badarinarayana, V., Audino, D.C., Rappsilber, J., Mann, M. & Denis, C.L.  
 1007 1998. The NOT proteins are part of the CCR4 transcriptional complex and affect gene  
 1008 expression both positively and negatively. *EMBO J.* **17**: 1096-1106.  
 1009  
 1010 Lushchak, V.I. & Gospodaryov, D.V. 2005. Catalases protect cellular proteins from  
 1011 oxidative modification in *Saccharomyces cerevisiae*. *Cell Biol. Int.* **29**: 187-192.



1012  
1013 Maathuis, F.J.M. & Amtmann, A. 1999. K<sup>+</sup> Nutrition and Na<sup>+</sup> Toxicity : The Basis of  
1014 Cellular K<sup>+</sup>/Na<sup>+</sup> Ratios. *Annals of Bot.* **84**: 123-133.  
1015  
1016 Mai, B. & Breeden, L. 1997. Xbp1, a Stress-Induced Transcriptional Repressor of the  
1017 *Saccharomyces cerevisiae* Swi4/Mbp1 Family. *Mol. Cell. Biol.* **17**: 6491-6501.  
1018  
1019 Manolio, T.A., Collins, F.S., Cox, N.J., Goldstein, D.B., Hindorff, L.A., Hunter, D.J. *et al.*  
1020 2009. Finding the missing heritability of complex diseases. *Nature* **461**: 747-753.  
1021  
1022 Margulies, M., Egholm, M., Altman, W.E., Attiya, S., Bader, J.S., Bemben, L.A. *et al.*  
1023 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature*  
1024 **437**: 376-380.  
1025  
1026 Martinez-Pastor, M.T., Marchler, G., Schuller, C., Marchler-Bauer, A., Ruis, H. &  
1027 Estruch, F. 1996. The *Saccharomyces cerevisiae* zinc finger proteins Msn2p and Msn4p  
1028 are required for transcriptional induction through the stress-response element (STRE).  
1029 *EMBO J.* **15**: 2227-2235.  
1030  
1031 Mendizabal, I., Rios, G., Mulet, J.M., Serrano, R. & de Larrinoa, I.F. 1998. Yeast  
1032 putative transcription factors involved in salt tolerance. *FEBS Lett.* **425**: 323-328.  
1033  
1034 Mendoza, I., Rubioli, F., Rodriguez-Navarro, A. & Pardo, J.M. 1994. The Protein  
1035 Phosphatase Calcineurin Is Essential for NaCl Tolerance of *Saccharomyces cerevisiae*.  
1036 *J. Biol. Chem.* **269**: 8792-8796.  
1037  
1038 Mersman, D.P., Du, H.N., Fingerman, I.M., South, P.F. & Briggs, S.D. 2009.  
1039 Polyubiquitination of the demethylase Jhd2 controls histone methylation and gene  
1040 expression. *Genes Dev.* **23**: 951-962.  
1041  
1042 Morgan, L.W. & Feldman, J.F. 2001. Epistatic and Synergistic Interactions Between  
1043 Circadian Clock Mutations in *Neurospora crassa*. *Genetics* **159**: 537-543.  
1044  
1045 Moskva, E., Schuller, C., Maurer, C.T.C., Mager, W.H. & Ruis, H. 1998. A Search in  
1046 the Genome of *Saccharomyces cerevisiae* for Genes Regulated via Stress Response  
1047 Elements. *Yeast* **14**: 1041-1050.  
1048  
1049 Mulet, J.M., Leube, M.P., Kron, S.J., Rios, G., Fink, G.R. & Serrano, R. 1999. A novel  
1050 mechanism of ion homeostasis and salt tolerance in yeast: the Hal4 and Hal5 protein  
1051 kinases modulate the Trk1-Trk2 potassium transporter. *Mol. Cell. Biol.* **19**: 3328-3337.  
1052  
1053 Mukhopadhyay, A., He, Z., Alm, E.J., Arkin, A.P., Baidoo, E.E., Borglin, S.C. *et al.* 2006:  
1054 Salt Stress in *Desulfovibrio vulgaris* Hildenborough: an Integrated Genomics Approach.  
1055 *J. Bacteriol.* **188**: 4068-4078.

1056  
1057 Murguía, J.R., Bellés, J.M. & Serrano, R. 1996. The yeast HAL2 nucleotidase is an in  
1058 vivo target of salt toxicity. *J. Biol. Chem.* **271**: 29029-29033.  
1059  
1060 Neiman, A.M. 1993. Conservation and reiteration of a kinase cascade. *Trends Genet.* **9**:  
1061 390-394.  
1062  
1063 Neiman, A.M., Stevenson, B.J., Xu, H-P., Sprague, Jr. G.F., Herskowitz, I., Wigler, M. et  
1064 al. 1993. Functional Homology of Protein Kinases Required for Sexual Differentiation in  
1065 *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae* Suggests a Conserved  
1066 Signal Transduction Module in Eukaryotic Organisms. *Mol. Biol. Cell* **4**: 107-120.  
1067  
1068 Noggle, R. 1946. The physiology of polyploidy in plants. I. Review of the literature.  
1069 *Lloydia* **9**; 153-173.  
1070  
1071 Nishida, E. & Gotoh, Y. 1993. The MAP kinase cascade is essential for diverse signal  
1072 transduction pathways. *Trends Biochem. Sci.* **18**: 128-131.  
1073  
1074 Ostrander, D.B. & Gorman, J.A. 1999. The Extracellular Domain of the *Saccharomyces*  
1075 *cerevisiae* Sln1p Membrane Osmolarity Sensor Is Necessary for Kinase Activity. *J.*  
1076 *Bacteriol.* **181**: 2527-2534.  
1077  
1078 Panaretou, B. & Piper, P.W. 1992. The plasma membrane of yeast acquires a novel  
1079 heat-shock protein (hsp30) and displays a decline in proton-pumping ATPase levels in  
1080 response to both heat shock and the entry to stationary phase. *Eur. J. Biochem.* **206**:  
1081 635-640.  
1082  
1083 Panasenko, O., Landrieux, E., Feuermann, M., Finka, A., Paquet, N. & Collart, M.A.  
1084 2006. The yeast Ccr4-Not complex controls ubiquitination of the nascent-associated  
1085 polypeptide (NAC-EGD) complex. *J. Biol. Chem.* **281**: 31389-31398.  
1086  
1087 Pardo, J.M., Reddy, M.P., Yang, S., Maggio, A., Huh, G-H., Matsumoto, T. et al. 1998.  
1088 Stress signaling through Ca<sup>2+</sup>/calmodulin-dependent protein phosphatase calcineurin  
1089 mediates salt adaptation in plants. *Proc. Natl. Acad. Sci. USA* **95**: 9681-9686.  
1090  
1091 Park, J.H., Wacholder, S., Gail, M.H., Peters, U., Jacobs, K.B., Chanock, S.J. et al.  
1092 2010. Estimation of effect size distribution from genome-wide association studies and  
1093 implications for future discoveries. *Nat. Genet.* **42**: 570-575.  
1094  
1095 Pelech, S.L. & Sanghera, J.S. 1992. Mitogen-activated protein kinases: versatile  
1096 transducers for cell signaling. *Trends Biochem. Sci.* **17**: 233-238.  
1097  
1098 Philippe, N., Pelosi, L., Lenski, R.E. & Schneider, D. 2009. Evolution of Penicillin-  
1099 Binding Protein 2 Concentration and Cell Shape during a Long-Term Experiment with

1100 *Escherichia coli*. *J. Bacteriol.* **191**: 909-921.  
 1101  
 1102 Phillips, P.C. 2008. Epistasis - the essential role of gene interactions in the structure  
 1103 and evolution of genetic systems. *Nat. Rev. Genet.* **9**: 855-867.  
 1104  
 1105 Piper, P.W., Ortiz-Calderon, C., Holyoak, C., Coote, P. & Cole, M. 1997. Hsp30, the  
 1106 integral plasma membrane heat shock protein of *Saccharomyces cerevisiae*, is a stress-  
 1107 inducible regulator of plasma membrane H(+)-ATPase. *Cell Stress Chaperones* **2**: 12-  
 1108 24.  
 1109  
 1110 Piper, P.W., Talreja, K., Panaretou, B., Moradas-Ferreira, P., Byrne, K., Praekelt, U.M.  
 1111 *et al.* 1994. Induction of major heat-shock proteins of *Saccharomyces cerevisiae*,  
 1112 including plasma membrane Hsp30, by ethanol levels above a critical threshold.  
 1113 *Microbiology* **140**: 3031-3038.  
 1114  
 1115 Posas, F., Chambers, J.R., Heyman, J.A., Hoeffler, J.P., de Nadal, E. & Arino, J. 2000.  
 1116 The Transcriptional Response of Yeast to Saline Stress. *J. Biol. Chem.* **275**: 17249-  
 1117 17255.  
 1118  
 1119 Quintero, F.J., Blatt, M.R. & Pardo, J.M. 2000. Functional conservation between yeast  
 1120 and plant endosomal Na<sup>+</sup>/H<sup>+</sup> antiporters. *FEBS Lett.* **471**: 224-228.  
 1121  
 1122 Quintero, F.J., Ohta, M., Shi, H., Zhu, J-K. & Pardo, J.M. 2002. Reconstitution in yeast  
 1123 of the *Arabidopsis* SOS signaling pathway for Na<sup>+</sup> homeostasis. *Proc. Natl. Acad. Sci.*  
 1124 *USA* **99**: 9061-9066.  
 1125  
 1126 Rancati, G., Pavelka, N., Fleharty, B., Noll, A., Trimble, R., Walton, K. *et al.* 2008.  
 1127 Aneuploidy underlies rapid adaptive evolution of yeast cells deprived of a conserved  
 1128 cytokinesis motor. *Cell* **135**: 879-893.  
 1129  
 1130 Reiser, V., Salah, S.M. & Ammerer, G. 2000. Polarized localization of yeast Pbs2  
 1131 depends on osmostress, the membrane protein Sho1 and Cdc42. *Nat. Cell. Biol.* **2**: 620-  
 1132 627.  
 1133  
 1134 Rep, M., Reiser, V., Gartner, U., Thevelein, J.M., Hohmann, S., Ammerer, G. *et al.*  
 1135 1999. Osmotic Stress-Induced Gene Expression in *Saccharomyces cerevisiae* Requires  
 1136 Msn1p and the Novel Nuclear Factor Hot1p. *Mol. Cell. Biol.* **19**: 5474-5485.  
 1137  
 1138 Rep, M., Krantz, M., Thevelein, J.M. & Hohmann, S. 2000. The transcriptional response  
 1139 of *Saccharomyces cerevisiae* to osmotic shock. *J. Biol. Chem.* **275**: 8290-8300.  
 1140  
 1141 Ringrose, L. & Paro, R. 2004. Epigenetic regulation of cellular memory by the Polycomb  
 1142 and Trithorax group proteins. *Annu. Rev. Genet.* **38**: 413-443.  
 1143

1144 Rios, G., Ferrando, A. & Serrano, R. 1997. Mechanisms of salt tolerance conferred by  
 1145 overexpression of the HAL1 gene in *Saccharomyces cerevisiae*. *Yeast* **13**: 515-528.  
 1146  
 1147 Rudolph, B., Gebendorfer, K.M., Buchner, J. & Winter, J. 2010. Evolution of *Escherichia*  
 1148 *coli* for Growth at High Temperatures. *J. Biol. Chem.* **285**: 19029-19034.  
 1149  
 1150 Saito, H. & Tatebayashi, K. 2004. Regulation of the Osmoregulatory HOG MAPK  
 1151 Cascade in Yeast. *J. Biochem.* **136**: 267-272.  
 1152  
 1153 Saleh, B., Allario, T., Dambier, D., Ollitrault, P. & Morillon, R. 2008. Tetraploid citrus  
 1154 rootstocks are more tolerant to salt stress than diploid. *C. R. Biologies*, **331**: 703-710.  
 1155  
 1156 Samani, P. & Bell, G. 2010. Adaptation of experimental yeast populations to stressful  
 1157 conditions in relation to population size. *J. Evol. Biol.* **23**: 791-796.  
 1158  
 1159 Sambrook, J. & Russell, D.W. 2001. *Molecular Cloning: A laboratory Manual*, 3rd edn.  
 1160 Cold Spring Harbor Laboratory Press, New York.  
 1161  
 1162 Sanchez, I., Hughes, R.T., Mayer, B.J., Yee, K., Woodgett, J.R. & Avruch, J. *et al.* 1994.  
 1163 Role of SAPK/ERK kinase-1 in the stress-activated pathway regulating transcription  
 1164 factor c-Jun. *Nature* **372**: 794-798.  
 1165  
 1166 Sanders, D. 2000. Plant biology: The salty tale of *Arabidopsis*. *Curr. Biol.* **10**: R486-  
 1167 R488  
 1168  
 1169 Schmitt, A.P. & McEntee, K. 1996. Msn2p, a zinc finger DNA-binding protein, is the  
 1170 transcriptional activator of the multistress response in *Saccharomyces cerevisiae*. *Proc.*  
 1171 *Natl. Acad. Sci. USA* **93**: 5777-5782.  
 1172  
 1173 Schuller, C., Brewster, J.L., Alexander, M.R., Gustin, M.C. & Ruis, H. 1994. The HOG  
 1174 pathway controls osmotic regulation of transcription via the stress response element  
 1175 (STRE) of the *Saccharomyces cerevisiae* CTT1 gene. *EMBO J.* **13**: 4382-4389.  
 1176  
 1177 Serrano, R. 1996. Salt tolerance in plants and microorganisms: toxicity targets and  
 1178 defense responses. *Int. Rev. Cytol.* **165**: 1-52.  
 1179  
 1180 Serrano, R., Mulet, J.M., Rios, G., Marquez, J.A., de Larrinoa, I.F., Leube, M.P. *et al.*  
 1181 1999. A glimpse of the mechanisms of ion homeostasis during salt stress. *J. Exp. Bot.*  
 1182 **50**: 1023-1036.  
 1183  
 1184 Stanek, M.T., Cooper, T.F. & Lenski, R.E. 2009. Identification and dynamics of a  
 1185 beneficial mutation in a long-term evolution experiment with *Escherichia coli*. *BMC Evol.*  
 1186 *Biol.* **9**:302.  
 1187

1188 Stebbins, G. L. 1950. Variation and evolution in plants. *Columbia University Press*, New  
1189 York.  
1190  
1191 Tal, M., & Gardi, I. 1976. Physiology of polyploid plants: water balance in autotetraploid  
1192 and diploid tomato under low and high salinity. *Physiol. Plantarum* **38**: 257-261.  
1193  
1194 Teige, M., Scheikl, E., Eulgem, T., Doczi, R., Ichimura, K., Shinozaki, K. *et al.* 2004. The  
1195 MKK2 Pathway Mediates Cold and Salt Stress Signaling in *Arabidopsis*. *Mol. Cell* **15**:  
1196 141-152.  
1197  
1198 Tsuda, L., Inoue, Y.H., Yoo, M.A., Mizuno, M., Hata, M., Lim, Y.M. *et al.* 1993. A protein  
1199 kinase similar to MAP kinase activator acts downstream of the raf kinase in *Drosophila*.  
1200 *Cell* **72**: 407-414.  
1201  
1202 Tsujimoto, Y., Izawa, S. & Inoue, Y. 2000. Cooperative Regulation of DOG2, Encoding  
1203 2-Deoxyglucose-6-Phosphate Phosphatase, by Snf1 Kinase and the High-Osmolarity  
1204 Glycerol-Mitogen-Activated Protein Kinase Cascade in Stress Responses of  
1205 *Saccharomyces cerevisiae*. *J. Bacteriol.* **182**: 5121-5126.  
1206  
1207 Trainotti, N. & Stambuk, B. U. 2001. NaCl stress inhibits maltose fermentation by  
1208 *Saccharomyces cerevisiae*. *Biotech. Lett.* **23**: 1703-1707.  
1209  
1210 Turner, B.M. 2002. Cellular Memory and the Histone Code. *Cell* **111**: 285-291.  
1211  
1212 Velicer, G.J., Raddatz, G., Keller, H., Deiss, S., Lanz, C., Dinkelacker, I. *et al.* 2006.  
1213 Comprehensive mutation identification in an evolved bacterial cooperator and its  
1214 cheating ancestor. *Proc. Natl. Acad. Sci. USA* **103**: 8107-8112.  
1215  
1216 Vijaranakul, U., Nadakavukaren, M.J., de Jonge, B.L., Wilkinson, B.J. & Jayaswal, R.K.  
1217 1995. Increased Cell Size and Shortened Peptidoglycan Interpeptide Bridge of NaCl-  
1218 Stressed *Staphylococcus aureus* and Their Reversal by Glycine Betaine. *J. Bacteriol.*  
1219 **177**: 5116-5121.  
1220  
1221 Visscher, P.M. 2008. Sizing up human height variation. *Nat. Genet.* **40**: 489-490.  
1222  
1223 Visscher, P.M. & Montgomery, G.W. 2009. Genome-wide Association Studies and  
1224 Human Disease: From Trickle to Flood. *J. Am. Med. Assoc.* **302**: 2028-2029.  
1225  
1226 Warringer, J., Ericson, E., Fernandez, L., Nerman, O. & Blomberg, A. 2003. High-  
1227 resolution yeast phenomics resolves different physiological features in the saline  
1228 response. *Proc. Natl. Acad. Sci. USA* **100**: 15724-15729.  
1229  
1230 Waskiewicz, A.J. & Cooper, J.A. 1995. Mitogen and stress response pathways: MAP  
1231 kinase cascades and phosphatase regulation in mammals and yeast. *Curr. Opin. Cell*

1232 *Biol.* **7**:798-805.

1233

1234 Wittkopp, P.J., Haerum, B.K. & Clark, A.G. 2004. Evolutionary changes in cis and trans

1235 gene regulation. *Nature* **430**: 85-88.

1236

1237 Wu, C-Y., Rolfe, P.A., Gifford, D.K. & Fink, G.R. 2010. Control of Transcription by Cell

1238 Size. *PLoS Biol* **8**: e1000523.

1239

1240 Wu, X., Hart, H., Cheng, C., Roach, P.J. & Tatchell, K. 2001. Characterization of Gac1p,

1241 a regulatory subunit of protein phosphatase type I involved in glycogen accumulation in

1242 *Saccharomyces cerevisiae*. *Mol. Genet. Genomics* **265**: 622-635.

1243

1244 Yale, J. & Bohnert, H.J. 2001. Transcript Expression in *Saccharomyces cerevisiae* at

1245 High Salinity. *J. Biol. Chem.* **276**: 15996-16007.

1246

1247 Yan, M., Dai, T., Deak, J.C., Kyriakis, J.M., Zon, L.I., Woodgett, J.R. *et al.* 1994.

1248 Activation of stress-activated protein kinase by MEKK1 phosphorylation of its activator

1249 SEK1. *Nature* **372**: 798-800.

1250

1251 Yang, S.X., Zhao, Y.X., Zhang, Q., He, Y.K., Zhang, H. & Luo, D. 2001. HAL1 mediate

1252 salt adaptation in *Arabidopsis thaliana*. *Cell Res.* **11**: 142-148.

1253

1254 Yang, J., Benyamin, B., McEvoy, B.P., Gordon, S., Henders, A.K., Nyholt, D.R. *et al.*

1255 2010. Common SNPs explain a large proportion of the heritability for human height. *Nat.*

1256 *Genet.* **42**: 565-569.

1257

1258 Yashar, B.M., Kelley, C., Yee, K., Errede, B. & Zon, L.I. 1993. Novel Members of the

1259 Mitogen-Activated Protein Kinase Activator Family in *Xenopus laevis*. *Mol. Cell. Biol.* **13**:

1260 5738-5748.

1261

1262 Zacharioudakis, I., Gligoris, T. & Tzamarias, D. 2007. A Yeast Catabolic Enzyme

1263 Controls Transcriptional Memory. *Curr. Biol.* **17**: 2041–2046.

1264

1265 Zamanillo, D., Hunt, T. & Nebreda, A.R. 1994. A novel kinase cascade triggered by

1266 chemical stress and heat shock which stimulates MAP kinase-activated protein kinase-2

1267 and phosphorylation of the small heat shock proteins. *Cell* **78**: 1027-1037.

1268

1269 Zeggini, E., Scott, L.J., Saxena, R., Voight, B.F., Marchini, J.L., Hu, T. *et al.* 2008. Meta-

1270 analysis of genome-wide association data and large-scale replication identifies

1271 additional susceptibility loci for type 2 diabetes. *Nat. Genet.* **40**: 638-645.

1272

1273 Zeyl, C., Vanderford, T. & Carter, M. 2003. An Evolutionary Advantage of Haploidy in

1274 Large Yeast Populations. *Science* **299**: 555-558.

1275

1276 Zhang, Y., Fatima, N. & Dufau, M.L. 2005. Coordinated Changes in DNA Methylation  
1277 and Histone Modifications Regulate Silencing/Derepression of Luteinizing Hormone  
1278 Receptor Gene Transcription. *Mol. Cell. Biol.* **25**: 7929-7939.

1279

1280 Zheng, D.Q., Wu, X.C., Tao, X.L., Wang, P.M., Li, P., Chi, X.Q., Li, Y.D., Yan, Q.F. &  
1281 Zhao, Y.H. 2011. Screening and construction of *Saccharomyces cerevisiae* strains with  
1282 improved multi-tolerance and bioethanol fermentation performance. *Bioresour. Technol.*  
1283 **102**:3020-3027.

1284

1285 Zhu, J-K. 2001. Plant salt tolerance. *Trends Plant Sci.* **6**: 66-71.

1286

1287 Zhu, J-K. 2002. Salt and Drought Stress Signal Transduction in Plants. *Annu. Rev.*  
1288 *Plant. Biol.* **53**: 247-273.

1289

1290 **Figures**

1291 **Figure 1 - (a-c) Viability of the ancestral strain and the evolved lines in medium**  
1292 **containing 0.5M NaCl.** Cells from the ancestral strain and the evolved lines from  
1293 overnight cultures were transferred into YPG medium. Cell samples from these cultures  
1294 were withdrawn after 16 hours (during exponential growth phase) as well as after 24  
1295 hours (stationary phase), and then diluted and plated. The plates were incubated at  
1296 30°C for five days and the number of colonies were counted. The relative viability of  
1297 both the ancestral strain and the evolved lines were estimated from the ratio of colonies  
1298 formed on agar plates with YPGN medium to that of plates with YPG medium. Relative  
1299 viabilities were measured for (a) the ancestral strain, (b) lines evolved in YPGN medium  
1300 (S lines), and (c) lines evolved in YPG medium (C lines). From the figures, it is clear that  
1301 NaCl does not affect the viability of the ancestral strain. The viabilities of the evolved  
1302 lines S1, S2 and S3 do not increase during the course of adaptation.

1303

1304 **Figure 2 - Fitness Assay of the evolved yeast lines.** Fitness assays were performed  
1305 by inoculating equal numbers of cells from an evolved line and from the GFP-tagged  
1306 reference yeast strain in the same vessel. The cells were grown for 24hrs, and the  
1307 percentage of non-GFP-tagged cells was counted using FACS. The ratio of the cell  
1308 numbers of the evolved lines to that of the ancestral strain gives the relative fitness of  
1309 the evolved lines. (a) Relative fitness of the three replicate yeast lines evolved in  
1310 medium with salt (S1,S2 and S3) in YPGN medium, and the three replicate yeast lines  
1311 evolved without NaCl (C1,C2,C3) measured in YPGN medium. (b) Relative fitness of  
1312 the lines S1, S2 and S3 in medium without NaCl. The lines evolved in salt show an  
1313 increase in growth rate by 8% to 12% in NaCl medium compared to the ancestral strain.  
1314 These lines also show a growth rate increase in medium without NaCl. However, this  
1315 increase in growth rate is consistently lower than the increase in NaCl medium. In  
1316 addition, the lines evolved in medium without NaCl show lower fitness in NaCl medium.  
1317 Taken together, it can be concluded that part of the adaptation in the salt evolved lines  
1318 is specific to NaCl.

1319

**Figure 3 - The genes showing similar (t-test, p-value<0.05, False Discovery Rate<10%) induction or repression by NaCl in the ancestral strain and in the evolved S lines** (a) Volcano plot, showing the p-values on the Y axis, and the fold expression change for these genes on the X axis. The labeled genes include genes associated with the stress response (e.g., *GPD1*, *SIP18*), genes with high level of expression change (e.g., *FMP48*, *NOG1*), and genes in both categories (e.g., *GRE1*). (b) Scatter plot for the fold-change of the common genes in the ancestral strain (X-axis) and in the evolved S lines (Y-axis). Although most of the common genes change expression to a similar extent in the ancestral strain and the S lines, a small number of genes change expression to a different extent.

**Figure 4 - Schematic diagram for the type of expression change that can occur as a result of evolutionary adaptation in the evolved S lines compared to the ancestral strain.** Each bar chart reflects the expression level of a hypothetical gene in the ancestral and evolved strain, in the absence and presence of salt. The scenarios that are depicted are as follows (a) No basal change, increased regulation (b) No basal change, increased repression (c) Basal increase in expression but no change in regulation (d) Basal decrease in expression but no change in regulation.

**Figure 5 - Genes with differential expression in the S lines compared to the ancestral strain ( $|Z\text{-score}|\geq 1.5$ ).** The plots show the level of expression change (vertical axis) and the Z-score (horizontal axis) for genes with (a) changed regulation, and (b) changed basal expression in the evolved lines. One should note that the basal change in expression is represented in terms of fold change, where the expression change is calculated as the ratio of the expressions in the evolved lines and the ancestral strain. The calculation of regulated expression change involves an additional subtraction of basal expression change and thus, can not be represented as fold change.

**Figure 6 - The five genes with the highest Z-score in each of four categories of expression change in the evolved lines.** Categories are (a) change in the level of induction, (b) change in the level of repression, (c) increase in basal expression, and (d) decrease in basal expression. The columns show, from left to right, the systematic gene name, the extent of expression change, the Z-score, the gene's common name, the type of expression change (in (a) and b) only), and a reference, if the gene had responded physiologically to hyperosmotic stress or saline stress in previous studies.

**Figure 7 - Different functional gene classes (Guldener *et al.*, 2005) are enriched or impoverished for genes that change expression in the salt evolved yeast lines (S lines), compared to the ancestral strain.** The vertical axis shows the functional classes of genes we consider, and the horizontal axis shows the fraction of genes in each category listed on the bottom of the figure. Asterisks (\*\*) indicate classes where the number of genes differs significantly from that expected by chance alone, as indicated by an exact binomial test.



1364

1365 **Figure 8 - Fitness analysis (using FACS) of the ancestral strain into which the**  
1366 **mutant *MOT2* gene of line S2 was introduced, and the corresponding wild-type**  
1367 ***MOT2* gene as a control.** (a) Fitness in medium with NaCl (b) Fitness in medium  
1368 without NaCl. The data show that the mutant *MOT2* gene increases the fitness in salt  
1369 medium as well as in medium without salt, and to similar extents.

1370

1371 **Figure 9 - Pulsed field gel electrophoresis (PFGE) analysis of the three replicate**  
1372 **salt evolved lines S1,S2 & S3 and the ancestral strain.** Two clones from each of the  
1373 lines S1,S2 and S3 were subjected to PFGE analysis along with the ancestral strain. (a)  
1374 The PFGE for the yeast chromosomes from the evolved lines S1, S2 and S3 and the  
1375 ancestral strain. The chromosomes are shown on the left side. M represents the yeast  
1376 chromosome PFG marker (NEB Catalog# N0345S). Only line S3 shows a new band  
1377 (red arrow) of size approximately 500kb. (b) PFGE analysis of the *NotI* digested  
1378 chromosomes from the lines S1,S2,S3 and the ancestral strain. M represents the Mid-  
1379 range PFG marker I from NEB (Catalog# N3551S), whose fragments range between 15  
1380 kb and 291 kb. Again, only line S3 shows a new band (red arrow) in the gel, which has  
1381 a size of approximately 180kb.

1382

1383 **Figure 10 - DNA content per yeast cell of the ancestral strain and the evolved**  
1384 **lines.**

1385 The ancestral strain and population samples of the evolved lines were grown for 24  
1386 hours at 30°C. For each of the cultures, the cell density was estimated. Genomic DNA  
1387 was isolated from a defined number of cells and DNA quantification was carried out with  
1388 a NanoDrop ND-1000 spectrophotometer. The DNA content per cell was then  
1389 calculated for each line. The figure suggests that all the evolved lines have become  
1390 massively aneuploid.

1391

1392 **Figure 11 - Cell size comparison between ancestral strain and one evolved line.**

1393 (a) Histogram of cell diameters of the ancestral strain and the S2 line, showing that the  
1394 salt evolved cells are bigger than the ancestral cells. (b) Microphotograph of cell  
1395 samples from the ancestral strain and the salt evolved lines.

1396

1397

1398

1399

1400

1401

1402

1403

1404


1405

1406



1407

1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451


Figure 1:




1452  
1453  
1454 Figure 2:



1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510 (b)



1528 Figure




4:

1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564

1565 Figure 5:  
1566 (a)

1567 Change in regulation: Expression change vs Z-score



1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602 Figure 6:

(a) Top five genes with change in induction level

Gene	Log2 (Expression Change)	Z-score	Common Name	Type of change	Previous studies
YOR348C	0.8821	2.088	PUT4	Increased induction	Posas et al., 2000; Rep et al., 2000; Yale & Bonhert, 2001
YHR053C	0.7017	1.8401	CUP1-1	Increased induction	Yale & Bonhert, 2001
YKL049C	0.554	1.8101	CSE4	New induction	Yale & Bonhert, 2001
YAR010C	0.9355	1.7493	---	New induction	Yale & Bonhert, 2001
YHR071W	0.5834	1.726	---	Increased induction	Yale & Bonhert, 2001

(b) Top five genes with change in repression level

Gene	Log2 (Expression Change)	Z-score	Common Name	Type of change	Previous studies
YMR189W	-1.0952	-2.9498	GCV2	New repression	Yale & Bonhert, 2001
YER091C	-0.9458	-2.6921	MET6	Increased repression	Rep et al., 2000
YNR053C	-1.0932	-2.6838	NOG2	Increased repression	---
YOR339C	-0.8424	-2.613	UBC11	Increased repression	---
YHR066W	-0.9644	-2.4702	SSF1	Increased repression	---


(c) Top five genes with increase in basal expression

Gene	Log2 (Fold Change)	Z-score	Common Name	Previous studies
YNL034W	1.3572	4.2657	---	---
YOR384W	1.2084	3.8216	FRE5	---
YOR049C	1.4081	3.5885	RSB1	---
YGL192W	1.5698	3.3941	IME4	---
YOR382W	1.8809	3.3935	FIT2	Yale & Bonhert, 2001

(d) Top five genes with decrease in basal expression

Gene	Log2 (Fold Change)	Z-score	Common Name	Previous studies
YKL178C	-1.2774	-3.9263	STE3	---
YML058W-A	-1.2118	-3.7933	HUG1	---
YJR078W	-1.2138	-3.5631	BNA2	---
YER011W	-1.0741	-3.361	TIR1	---
YDR281C	-1.0859	-2.789	PHM6	---


Figure 7:



1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661




1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676 Figure 8:




1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693

1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737


Figure 9:



1738  
1739  
1740 Figure 10:



1771  
1772  
1773  
1774  
1775  
1776  
1777 Figure 11:  
1778  
1779 (a)  
1780  
1781



1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796 (b)  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812

